



ADDITIONS TO THE TAIWAN EEL FAUNA WITH FIVE NEWLY RECORDED SPECIES OF MORAY EELS (ANGUILLIFORMES: MURAENIDAE), AND REDESCRIPTION OF A RARE SPECIES *Gymnothorax sagmacephalus*

Kar-Hoe Loh

Institute of Ocean and Earth Sciences, University of Malaya, Kuala Lumpur, Malaysia 50603

Kwang-Tsao Shao

Biodiversity Research Center, Academia Sinica, Nankang, Taiwan, R.O.C

Ving-Ching Chong

Institute of Ocean and Earth Sciences, University of Malaya, Kuala Lumpur, Malaysia 50603. Institute of Biological Sciences, University of Malaya, Kuala Lumpur, Malaysia 50603

Hong-Ming Chen

Department of Aquaculture, National Taiwan Ocean University, Keelung, Taiwan, R.O.C. 5 Center of Excellence for the Oceans, National Taiwan Ocean University, Keelung, Taiwan, R.O.C, hmchen@mail.ntou.edu.tw

Follow this and additional works at: <https://jmstt.ntou.edu.tw/journal>



Part of the [Aquaculture and Fisheries Commons](#)

Recommended Citation

Loh, Kar-Hoe; Shao, Kwang-Tsao; Chong, Ving-Ching; and Chen, Hong-Ming (2016) "ADDITIONS TO THE TAIWAN EEL FAUNA WITH FIVE NEWLY RECORDED SPECIES OF MORAY EELS (ANGUILLIFORMES: MURAENIDAE), AND REDESCRIPTION OF A RARE SPECIES *Gymnothorax sagmacephalus*," *Journal of Marine Science and Technology*: Vol. 24: Iss. 2, Article 19.

DOI: 10.6119/JMST-013-1227-1

Available at: <https://jmstt.ntou.edu.tw/journal/vol24/iss2/19>

This Research Article is brought to you for free and open access by Journal of Marine Science and Technology. It has been accepted for inclusion in Journal of Marine Science and Technology by an authorized editor of Journal of Marine Science and Technology.

ADDITIONS TO THE TAIWAN EEL FAUNA WITH FIVE NEWLY RECORDED SPECIES OF MORAY EELS (ANGUILLIFORMES: MURAENIDAE), AND REDESCRIPTION OF A RARE SPECIES *Gymnothorax sagmacephalus*

Acknowledgements

We are very grateful to Captain Jiunn-Shiun Chiou and Ms. Min-Chia Chiang, who helped us to collect these specimens, and two anonymous reviewers for improving the manuscript. These studies were originally supported by National Science Council "NSC 99-2621-B-001-008-MY3; 98-2631-H-019-001"; Fisheries Agency, Council of Agriculture grants "98AS8.5.2-FA-F1 (2-1)" to Hong-Ming Chen; and University Malaya short research fund "BK027-2011A". Continuing studies of morays have greatly benefitted from discussions with J. E. McCosker (CAS), D. G. Smith (USNM), and J. E. Randall (BPBM). The authors also acknowledge Prof. Chang-Tai Shih for his valuable comments and English editing support.

ADDITIONS TO THE TAIWAN EEL FAUNA WITH FIVE NEWLY RECORDED SPECIES OF MORAY EELS (ANGUILLIFORMES: MURAENIDAE), AND REDESCRIPTION OF A RARE SPECIES *Gymnothorax sagmacephalus*

Kar-Hoe Loh¹, Kwang-Tsao Shao², Ving-Ching Chong^{1, 3}, and Hong-Ming Chen^{4, 5}

Key words: Muraenidae, *Anarchias*, *Gymnothorax*, Taiwan.

ABSTRACT

Five rare moray species: *Anarchias cantonensis* (Schultz), *Gymnothorax dorsalis* Seale, *G. intesi* (Fourmanoir & Rivaton), *G. phasmatodes* (Smith) and *G. sagmacephalus* (Böhlke) belonging to the subfamilies Uropterygiinae and Muraeninae are reported for the first time in the surrounding waters off Taiwan. Diagnoses of these five species, including fresh coloration and dentition, are provided. A re-description of *G. sagmacephalus* is provided since its original description, based on the single holotype, was incomplete specifically due to its damaged tail. With the added 12 specimens, we provided the statistics of the morphological measurements and vertebral formulae, and expanded its geographical distribution from Japan southward to Taiwan.

I. INTRODUCTION

Moray eels live in tropical and subtropical coastal areas, belong to the diverse eel family Muraenidae, and comprise 16 genera and 198 species globally (Eschmeyer and Fong, 2013). The Taiwan moray eel fauna was comprehensively treated by Chen et al. (1994) who reported nine genera and 42 species. During our surveys from 2005 to 2009 under the research

programme, “Diversity, molecular phylogeny and reproductive ecology of the Anguilliformes fishes of Taiwan and the western Pacific”, numerous specimens of moray eels from the seas around Taiwan were collected. New species and new records of moray eels were identified among them (Chen and Loh, 2007; Chen et al., 2008; Loh et al., 2008; Shao et al., 2008; Loh et al., 2011; Loh et al., 2012). In this study, we report five muraenid species new to Taiwan and amend the muraenid fauna of Taiwan to composing 68 species in 13 genera and two subfamilies (Table 1).

The aim of this paper was to list the muraenid eels from Taiwan, to describe the five newly recorded moray species, and to redescribe *Gymnothorax sagmacephalus*. Böhlke (1997) described *G. sagmacephalus* based on the single type specimen, (collected from Tokyo Bay, Japan), with missing tail tip. This species was subsequently reported by Böhlke and Smith (2002) as a valid species, characterized by an elongate body, tannish brown coloration, and vertebral count MVF 7-77-172+. We redescribed this species based on Böhlke’s description of the holotype and 12 additional recently collected specimens. Color photographs of a fresh specimen are also provided.

II. MATERIALS AND METHODS

All moray specimens were collected either by longlines or using the ichthyocide rotenone, from the surrounding waters of Taiwan. The specimens were fixed with 10% formalin, then transferred to 70% ethanol solution for long-term preservation. The methods of measurements followed Böhlke and Randall (2000). Proportional measurements for the specimens were expressed as percentage of the total length (TL) or the head length (HL). Body depths were measured at the gill openings (DGO) and at the anus (DA); snout length was measured from the snout tip to the anterior margin of the eye; upper jaw (UJ) length was measured from the snout tip to the rictus; lower jaw (LJ) length was measured from the lower jaw tip to the

Paper submitted 04/19/13; revised 12/12/13; accepted 12/27/13. Author for correspondence: Hong-Ming Chen (e-mail: hmchen@mail.ntou.edu.tw).

¹ Institute of Ocean and Earth Sciences, University of Malaya, Kuala Lumpur, Malaysia 50603.

² Biodiversity Research Center, Academia Sinica, Nankang, Taiwan, R.O.C.

³ Institute of Biological Sciences, University of Malaya, Kuala Lumpur, Malaysia 50603.

⁴ Department of Aquaculture, National Taiwan Ocean University, Keelung, Taiwan, R.O.C.

⁵ Center of Excellence for the Oceans, National Taiwan Ocean University, Keelung, Taiwan, R.O.C.

Table 1. The taxonomic works of all muraenid species in Taiwan.

| No. | Genus | Scientific name | Chen et al., 1997 | Chen and Loh, 2007 | Chen et al., 2008 | Loh et al., 2008 | Shao et al., 2008 | Loh et al., 2011 | Loh et al., 2012 | This study |
|-----|----------------------|--------------------------------------|----------------------|-----------------------|----------------------|---------------------|----------------------|---------------------|---------------------|------------|
| 1 | <i>Echidna</i> | <i>Echidna nebulosa</i> | ○ | | | | ○ | | | |
| 2 | | <i>Echidna polyzona</i> | ○ | | | | ○ | | | |
| 3 | | <i>Echidna xanthospilos</i> | ◎ | | | | | | | |
| 4 | <i>Enchelycore</i> | <i>Enchelycore bayeri</i> | ◎ | | | | | | | |
| 5 | | <i>Enchelycore bikiniensis</i> | ◎ | | | | | | | |
| 6 | | <i>Enchelycore lichenosa</i> | ○ | | | | | | | |
| 7 | | <i>Enchelycore pardalis</i> | ○ | | | | ○ | | | |
| 8 | | <i>Enchelycore schismatorhynchus</i> | ○ | | | | | | | |
| 9 | <i>Enchelynassa</i> | <i>Enchelynassa canina</i> | | | | | | | ◎ | |
| 10 | <i>Gymnomuraena</i> | <i>Gymnomuraena zebra</i> | ○ | | | | ○ | | | |
| 11 | <i>Gymnothorax</i> | <i>Gymnothorax albimarginatus</i> | ◎ | | | | | | | |
| 12 | | <i>Gymnothorax berndti</i> | ○ | | | | | | | |
| 13 | | <i>Gymnothorax buroensis</i> | ○ | | | | ○ | | | |
| 14 | | <i>Gymnothorax chilospilus</i> | ○ | | | | ○ | | | |
| 15 | | <i>Gymnothorax chlamydatus</i> | ○ | | | | | | | |
| 16 | | <i>Gymnothorax dorsalis</i> | | | | | | | | ◎ |
| 17 | | <i>Gymnothorax elegans</i> | | | | | | | | |
| 18 | | <i>Gymnothorax eurostus</i> | ○ | | | | ○ | | | |
| 19 | | <i>Gymnothorax favagineus</i> | ○ | | | | | | | |
| 20 | | <i>Gymnothorax fimbriatus</i> | ○ | | | | ○ | | ◎ | |
| 21 | | <i>Gymnothorax flavimarginatus</i> | ○ | | | | ○ | | | |
| 22 | | <i>Gymnothorax formosus</i> | ◎ | | | | | | | |
| 23 | | <i>Gymnothorax fuscomaculatus</i> | | | | | | | ◎ | |
| 24 | | <i>Gymnothorax gracilicauda</i> | | | | | ◎ | | | |
| 25 | | <i>Gymnothorax hepaticus</i> | ○ | | | | | | | |
| 26 | | <i>Gymnothorax herrei</i> | ◎ | | | | ○ | | | |
| 27 | | <i>Gymnothorax intesi</i> | | | | | | | | ◎ |
| 28 | | <i>Gymnothorax javanicus</i> | ◎ | | | | ○ | | | |
| 29 | | <i>Gymnothorax kidako</i> | ○ | | | | ○ | | | |
| 30 | | <i>Gymnothorax prionodon</i> | ○ | | | | | | | |
| 31 | | <i>Gymnothorax margaritophorus</i> | ◎ | | | | ○ | | | |
| 32 | | <i>Gymnothorax melanosomatus</i> | | | | | | ● | | |
| 33 | | <i>Gymnothorax melatremus</i> | ◎ | | | | ○ | | | |
| 34 | | <i>Gymnothorax meleagris</i> | ○ | | | | ○ | | | |
| 35 | | <i>Gymnothorax minor</i> | | | | | ○ | | | |
| 36 | | <i>Gymnothorax monostigma</i> | | | | | ○ | | | |
| 37 | | <i>Gymnothorax neglectus</i> | ○ | | | | ○ | | | |
| 38 | | <i>Gymnothorax niphostigmus</i> | ● | | | | | | | |
| 39 | | <i>Gymnothorax nudivomer</i> | ◎ | | | | | | | |
| 40 | | <i>Gymnothorax phasmatodes</i> | | | | | | | | ◎ |
| 41 | | <i>Gymnothorax pictus</i> | ○ | | | | ○ | | | |
| 42 | | <i>Gymnothorax pindae</i> | ○ | | | | ○ | | | |
| 43 | | <i>Gymnothorax polyuranodon</i> | | | | | ◎ | | | |
| 44 | | <i>Gymnothorax pseudothysoideus</i> | ○ | | | | ○ | | | |
| 45 | | <i>Gymnothorax reevesii</i> | ◎ | | | | ○ | | | |
| 46 | | <i>Gymnothorax reticularis</i> | ○ | | | | ○ | | | |
| 47 | | <i>Gymnothorax richardsoni</i> | ○ | | | | ○ | | | |
| 48 | | <i>Gymnothorax rueppelliae</i> | ○ | | | | ○ | | | |
| 49 | | <i>Gymnothorax sagmacephalus</i> | | | | | | | | ◎ |
| 50 | | <i>Gymnothorax shaoi</i> | | ● | | | | | | |
| 51 | | <i>Gymnothorax taiwanensis</i> | | | ● | | | | | |
| 52 | | <i>Gymnothorax thyrsoides</i> | ○ | | | | ○ | | | |
| 53 | | <i>Gymnothorax undulatus</i> | ○ | | | | ○ | | | |
| 54 | | <i>Gymnothorax ypsilon</i> | ○ | | | | | | | |
| 55 | | <i>Gymnothorax zonipectis</i> | ◎ | | | | ○ | | | |
| 56 | <i>Pseudechidna</i> | <i>Pseudechidna brummeri</i> | ○ | | | | | | | |
| 57 | <i>Rhinomuraena</i> | <i>Rhinomuraena quaesita</i> | ○ | | | | ○ | | | |
| 58 | <i>Strophidon</i> | <i>Strophidon prolatus</i> | ○ | | | | ○ | | | |
| 59 | | <i>Strophidon sathete</i> | ◎ | | | | ○ | | | |
| 60 | <i>Anarchias</i> | <i>Anarchias allardicei</i> | ○ | | | | ○ | | | |
| 61 | | <i>Anarchias cantonensis</i> | | | | | | | | ◎ |
| 62 | <i>Channomuraena</i> | <i>Channomuraena vittata</i> | ◎ | | | | ○ | | | |
| 63 | <i>Cirrimaxilla</i> | <i>Cirrimaxilla formosa</i> | ● | | | | | | | |
| 64 | <i>Scuticaria</i> | <i>Scuticaria tigrina</i> | ◎ | | | | ○ | | | |
| 65 | | <i>Scuticaria marmorata</i> | | | | ○ | ○ | | | |
| 66 | <i>Uropterygius</i> | <i>Uropterygius macrocephalus</i> | ◎ | | | | ○ | | | |
| 67 | | <i>Uropterygius micropterus</i> | ◎ | | | | ○ | | | |
| 68 | | <i>Uropterygius oligospondylus</i> | | | | ● | | | | |

○ presence, ◎ new record, ● new species.

rictus. Counts for the vertebral formulae were taken from radiographs, as explained in Böhlke (1982) and Chen et al. (1994, 1997); the mean vertebral formula (MVF) gives the mean values for predorsal-preanal-total vertebrae counts: predorsal vertebrae (PDV); preanal vertebrae (PAV); total vertebrae (TV). Teeth counts following Hatooka (1986) were approximate and included sockets of missing teeth. Sexes of the specimens were determined by gross and histological examinations of the gonads. All newly recorded specimens were deposited in the collection of the Laboratory of Aquatic Ecology, Department of Aquaculture, National Taiwan Ocean University (TOU-AE). Specimens were also borrowed from the following institutions: the National Museum of Marine Science and Technology, Pisces collection (NMMSTP); the Fisheries Research Institute (FRIP); and the Museum of Research Center for Biodiversity (Institute of Zoology), Academia Sinica (ASIZP).

Comparative materials examined.

Anarchias allardicei – Four specimens (116-154 mm TL), NMMSMP 00885, Tungchi, Penghu County; NMMSMP 01013, Wanlitung, Pingtung County; TOU-AE 1780 Hsiogang, Tai-tung County; TOU-AE 4913, Jihuei, Chengkung, Taitung County.

Gymnothorax albimarginatus – Five specimens (675-1060 mm TL), all male, TOU-AE 0104, 1638, 1813, from Bisha fishes market, Keelung City; TOU-AE 1034, 4220, Changbin, Taitung County.

Gymnothorax neglectus – Three specimens (610-699 mm TL), ASIZP 056655, Hopingtao, Keelung City; NMMSTP 00956, Hopingtao, Keelung City; FRIP 20370 Chengkung, Taitung County.

Strophidon sathete – 12 specimens (526-1470 mm TL), male: TOU-AE 1868, 3458, Changbin, Taitung County; TOU-AE 3026, 4832, 5305, Daxi, Yilan County; female: TOU-AE 0628, Changbin, Taitung County; TOU-AE 3027, 3028, 3990, 4478, 4563, 5306, Daxi, Yilan County.

III. TAXONOMY

Uropterygiinae

Diagnosis

Dorsal and anal fins restricted to tail tip; teeth needle-like and pointed.

Type genus

Uropterygius Rüppell, 1838.

Anarchias Jordan & Starks in Jordan and Seale, 1906

Diagnosis

A supraorbital pore is adjacent to each posterior nostril, appearing as double-pore posterior nostrils.

Type species

Anarchias allardicei Jordan & Starks in Jordan and Seale, 1906.




Fig. 1. The newly recorded species, lateral view of body and head. A, *Anarchias cantonensis*, TOU-AE 4913 (93 mm TL); B, *Gymnothorax dorsalis*, TOU-AE 4834 (619 mm TL); C, *G. intesi*, TOU-AE 2774 (533 mm TL); D, *G. phasmatodes*, TOU-AE 0227 (382 mm TL); E, *G. sagmacephalus*, TOU-AE 0226 (464 mm TL).

Anarchias cantonensis (Schultz, 1943)
(Figs. 1A, 3A; Table. 2)

Uropterygius cantonensis Schultz, 1943: 27 (Type locality: Phoenix Islands, southern central Pacific).

Anarchias cantonensis (Schultz, 1943), Böhlke and McCosker, 2001: 73; Böhlke and Smith, 2002: 157; McCosker and Stewart, 2006: 86; Reece et al., 2010: 423-425.

1. Material Examined

Three specimens: TOU-AE 4913, 4914, 4915, 97-154 mm TL, females, 31 May 2008, from Jihuei, Chengkung, Taitung County, all rotenoned by Ms. Min-Chia Chiang.

2. Diagnosis

Small sized, body pale with a network of wide brownish lines in irregular vertical reticulated bars, fins at tip of tail yellow (Fig. 1A). Anterior nostril in a short tube, posterior nostril above center of eye, a supraorbital pore adjacent to posterior nostril on both sides, appearing as double-pore posterior nostrils. Tail length 1.9-2.0 (1.9 ± 0.1), trunk length 2.8-2.9 (2.8 ± 0.1), depth at gill opening 18.5-22.2 (20.8 ± 2.0), body depth at anus 19.2-21.2 (20.0 ± 1.1), head length 7.5-8.1 (7.7 ± 0.3), all in TL. Length of upper jaw 2.5-3.1 (2.8 ± 0.3), length of lower jaw 2.5-3.1 (2.8 ± 0.3), interorbital width 6.4-9.6 (7.7 ± 1.7), snout length 5.7-6.4 (6.1 ± 0.4), eye

Table 2. Total lengths, proportions as percent of total length or head length, counts of teeth and vertebrae among the five newly recorded muraenid species from Taiwan.

| Species | <i>Anarchias cantonensis</i> | | <i>Gymnothorax dorsalis</i> | | <i>Gymnothorax intesi</i> | | <i>Gymnothorax phasmatodes</i> | | <i>Gymnothorax sagmacephalus</i> | |
|--|------------------------------|------------|-----------------------------|------------|---------------------------|------------|--------------------------------|------------|----------------------------------|------------|
| | Number of Specimens | | 3 | | 7 | | 6 | | 12 | |
| | Range | Mean ± SD | Range | Mean ± SD | Range | Mean ± SD | Range | Mean ± SD | Range | Mean ± SD |
| Total Length (mm) | 97-154 | | 415-1050 | | 343-652 | | 268-382 | | 375-512 | |
| Proportions as percent of total length | | | | | | | | | | |
| Preanal length | 47.5-48.7 | 48.2 ± 0.6 | 41.0-44.3 | 42.9 ± 1.7 | 50.0-50.0 | 50.3 ± 0.3 | 50.4-54.5 | 52.3 ± 1.9 | 50.4-52.8 | 51.2 ± 0.8 |
| Tail length | 51.3-52.6 | 51.8 ± 0.7 | 55.7-59.1 | 57.1 ± 1.7 | 49.1-50.0 | 49.7 ± 0.3 | 45.5-49.6 | 47.7 ± 1.9 | 47.2-49.6 | 48.8 ± 0.8 |
| Trunk length | 34.3-36.4 | 35.3 ± 1.1 | 31.4-33.6 | 32.6 ± 1.1 | 39.1-40.1 | 39.5 ± 0.4 | 40.9-44.6 | 43.0 ± 1.6 | 39.7-43.6 | 41.3 ± 1.1 |
| Depth at gill opening | 4.5-5.4 | 4.8 ± 0.5 | 3.5-4.0 | 3.7 ± 0.3 | 4.9-7.2 | 6.1 ± 0.8 | 2.7-3.6 | 3.1 ± 0.5 | 2.9-3.8 | 3.4 ± 0.3 |
| Depth at anus | 4.7-5.2 | 5.0 ± 0.3 | 3.2-3.6 | 3.4 ± 0.2 | 3.7-4.8 | 4.2 ± 0.4 | 1.7-3.2 | 2.5 ± 0.6 | 2.2-3.6 | 2.8 ± 0.3 |
| Predorsal length | -- | -- | 8.1-9.6 | 8.8 ± 0.7 | 7.5-10.8 | 9.2 ± 1.3 | 6.2-7.5 | 7.0 ± 0.6 | 6.4-10.0 | 8.1 ± 1.0 |
| Head length | 12.3-13.4 | 12.9 ± 0.6 | 9.5-11.1 | 10.4 ± 0.8 | 10.6-12.0 | 11.2 ± 0.6 | 9.6-10.1 | 9.8 ± 0.2 | 9.7-11.5 | 10.6 ± 0.6 |
| Proportions as percent of head length | | | | | | | | | | |
| Upper jaw | 32.2-39.9 | 35.7 ± 3.9 | 32.0-36.7 | 34.2 ± 2.3 | 39.2-43.2 | 41.8 ± 1.5 | 35.2-40.4 | 35.9 ± 2.8 | 34.0-45.3 | 38.2 ± 3.0 |
| Lower jaw | 31.9-39.8 | 35.6 ± 4.0 | 30.6-35.9 | 33.4 ± 2.6 | 38.2-57.4 | 43.2 ± 6.5 | 32.6-40.4 | 34.7 ± 3.3 | 32.2-44.9 | 37.4 ± 3.3 |
| Interorbital width | 10.5-15.6 | 13.4 ± 2.6 | 9.9-13.7 | 12.0 ± 1.9 | 11.2-17.4 | 14.1 ± 2.4 | 10.9-17.3 | 13.8 ± 3.2 | 12.2-16.7 | 14.4 ± 1.4 |
| Snout length | 15.7-17.6 | 16.5 ± 1.0 | 12.5-13.4 | 13.0 ± 0.5 | 15.5-20.2 | 18.5 ± 1.6 | 17.1-19.1 | 17.9 ± 1.0 | 15.5-19.7 | 18.0 ± 1.2 |
| Eye diameter | 7.3-11.3 | 9.2 ± 2.0 | 5.4-5.9 | 5.7 ± 0.3 | 9.4-10.8 | 10.2 ± 0.5 | 8.3-11.2 | 9.9 ± 1.5 | 9.2-11.8 | 10.3 ± 0.8 |
| Teeth | | | | | | | | | | |
| Premaxillary | 10-13 | | 5-7 | | 11-14 | | 6-7 | | 5-7 | |
| Median premaxillary | 9 | | 3 | | 0-1 | | 0-3 | | 0-3 | |
| Maxillary-inner | 2-12 | | 2-5 | | - | | - | | - | |
| Maxillary-outer | 24-34 | | 15-18 | | 16-19 | | 7-12 | | 6-10 | |
| Vomerine | 6-10 | | 3-7 | | 11-20 | | 4-9 | | 3-8 | |
| Dentary-inner | 5-15 | | 2-5 | | - | | - | | - | |
| Dentary-outer | 32-42 | | 18-23 | | 26-40 | | 12-20 | | 9-18 | |
| Vertebrae | | | | | | | | | | |
| Predorsal | 86-93 | 90 | 8-9 | 9 | 5-6 | 6 | 5-6 | 6 | 6-7 | 6 |
| Preanal | 90-95 | 92 | 66-69 | 68 | 65-70 | 67 | 73-77 | 75 | 74-78 | 76 |
| Total | 100-105 | 102 | 164-167 | 166 | 149-154 | 151 | 165-168 | 166 | 170-176 | 173 |

diameter 8.8-13.8 (11.3 ± 2.5), all in HL. Premaxillary teeth 10-13 each side, median premaxillary teeth 9, maxillary teeth biserial about 2-12 inner row and 24-34 outer row; vomerine teeth uniserial 6-10, dentary teeth biserial about 5-15 inner row and 32-42 outer row (Fig. 3A). Predorsal vertebrae 86-93, preanal vertebrae 90-95, total vertebrae 100-105; mean vertebral formula 90-92-102.

3. Distribution

Known from the western Pacific, including the South China Sea and seas around Taiwan.

4. Remarks

Species of *Anarchias* are readily identified by a posterior nostril contiguous with an enlarged supraorbital pore on each side, appearing as double-pore posterior nostrils. *A. cantonensis* is distinguished from its congener *A. allardicei*, by the head length 12.9 (vs. 10.0) % of TL; snout length 16.5 (vs. 13.2) %, eye diameter 9.2 (vs. 7.6) %, interorbital width 13.4 (vs. 9.2) %, all of HL. Böhlke and Smith (2002) reported its MVF 91-93-103, similar to MVF 91-93-104 in Böhlke and McCosker (2011); both MVFs were similar to that of the present study (MVF 90-92-102). The total vertebrae of *A. cantonensis* reported by Reece et al. (2010) were more widely ranged (98-108) than ours (100-105).

Muraeninae

Diagnosis

Dorsal-fin origin anterior to or near gill opening; and anal-fin origin just posterior to anus.

Type genus

Muraena Linnaeus 1758.

Gymnothorax Bloch, 1795

Diagnosis

Mouth closing completely; jaws slightly or not curved.

Type species

Gymnothorax reticularis Bloch, 1795

Gymnothorax dorsalis Seale, 1917
(Figs. 1B, 3B; Table 2)

Gymnothorax dorsalis Seale, 1917: 92 (Type locality: Hong Kong); Smith, 1994: 17; Böhlke, 1997: 96-97; Böhlke and Smith, 2002: 106.

1. Materials Examined

Three specimens: TOU-AE 0157, 1050 mm TL, 18 May

2002; TOU-AE 4834, 619 mm TL, 22 Jan. 2008; TOU-AE 5303, 415 mm TL, 19 Sept. 2008, all were females from Daxi, Yilan County.

2. Diagnosis

Body medium tan-brown, anus anterior to mid-body. Tail length 1.7-1.8 (1.8 ± 0.1), trunk length 3.0-3.2 (3.1 ± 0.1), body depth at gill opening 25.1-28.9 (27.1 ± 1.9), body depth at anus 28.1-31.2 (29.9 ± 1.6), predorsal length 10.5-12.4 (11.5 ± 1.0), head length 9.0-10.6 (9.7 ± 0.8), all in TL. Upper jaw length 2.7-3.1 (2.9 ± 0.2), lower jaw length 2.8-3.3 (3.0 ± 0.2), interorbital width 7.3-10.1 (8.5 ± 1.4), snout length 7.5-8.0 (7.7 ± 0.3), eye diameter 16.8-18.7 (17.5 ± 1.0), all in HL. Premaxillary teeth 5-7 each side, median premaxillary teeth 3; maxillary teeth biserial, 2-5 in inner row and 15-18 in outer row, vomerine teeth uniserial 3-7; dentary teeth biserial, 2-5 in inner row and 18-23 in outer row (Fig. 3B). Head pores typical, three supraorbital pores, four infraorbital pores, six mandibular pores.

Two small branchial pores, anterior to gill opening. Gill opening below mid-side. Predorsal vertebrae 8-9, preanal vertebrae 66-69, total vertebrae 164-167; mean vertebral formula 9-68-166.

3. Distribution

Known from the western Pacific, including Hong Kong, China, Taiwan and Vietnam.

4. Remarks

Gymnothorax sensu lato is the catch-all genus in the family Muraenidae. In the present paper we are temporarily maintaining the species *Gymnothorax dorsalis* in the genus *Gymnothorax*, with the understanding that its generic status may change when osteological and phylogenetic studies are undertaken. The overall external description of *G. dorsalis* is similar to that of *Strophidon sathete*; the anus is well anterior to midbody, the eye is closer to the snout tip than to the rictus, and the jaw teeth are biserial. We believe that it will be moved to the genus *Strophidon* in the future when a revision is available.

However, *Gymnothorax dorsalis* is still distinguished from *Strophidon sathete*, by the following features: lesser PAV and TV value of MVF 9-68-166 (vs. 9-81-194); greater body depth at gill opening 3.5-4.0 (vs. 1.8-2.9) % of total length.

Gymnothorax intesi (Fourmanoir & Rivaton, 1979)
(Figs. 1C, 3C; Table 2)

Lycodontis intesi Fourmanoir and Rivaton, 1979: 426 (Type locality: Loyalty Islands, South western Pacific).

Gymnothorax intesi (Fourmanoir et Rivaton, 1979), Böhlke and McCosker, 2001: 80; Böhlke and Smith 2002: 116.

1. Materials Examined

Seven specimens: TOU-AE 2774, 533 mm TL, female, 05 Aug. 2006; TOU-AE 4407, 4124, 2 specimens, 454-652 mm

TL, females, 12 Feb. 2007; TOU-AE 2773, 2775, 2776, 3 specimens, 343-502 mm TL, males, 05 Aug. 2006; TOU-AE 4128, 613 mm TL, male, 05 Sept. 2006, all were from Taitung County, longline, Captain Jiunn-Shiun Chiou.

2. Diagnosis

An elongate moray, body light brown, with yellow-green mucus when fresh. Head dusky or reticulated, body with irregular pale spots; fins with pale or white margins. Preanal length 1.9-2.0 (1.9 ± 0.1), tail length 2.0-2.1 (2.0 ± 0.1), trunk length 2.5-2.6 (2.5 ± 0.1), body depth at gill opening 13.9-20.5 (16.7 ± 2.4), body depth at anus 20.8-27.0 (24.1 ± 2.0), predorsal length 9.3-13.4 (11.1 ± 1.6), head length 8.4-9.5 (9.0 ± 0.4), all in TL. Upper jaw length 2.3-2.6 (2.4 ± 0.1), lower jaw length 2.4-2.6 (2.5 ± 0.1), interorbital width 5.7-8.9 (7.3 ± 1.2), snout length 4.9-6.5 (5.4 ± 0.5), eye diameter 9.3-10.7 (9.9 ± 0.5), all in HL. Pre-maxillary teeth 11-14 each side, median premaxillary teeth 0-1; maxillary teeth uniserial, about 16-19, vomerine teeth uniserial 11-20, dentary teeth uniserial about 26-40 each side (Fig. 3C). Head pores typical, two branchial pores. Predorsal vertebrae 5-6, preanal vertebrae 65-70, total vertebrae 149-154; mean vertebral formula 6-67-151.

3. Distribution

Known from the Indian Ocean; central and western Pacific, including Australia, New Caledonia, Japan and Taiwan.

4. Remarks

Gymnothorax intesi is distinguished from its congener *G. neglectus*, by the following features: more total vertebrae 149-151 (vs. 138-142); smaller body depth at gill opening 4.9-7.2 (vs. 7.5-9.4) % of total length; and longer snout length 15.5-20.2 (vs. 13.3-13.9) % of head length.

Gymnothorax phasmatodes (Smith, 1962)
(Figs. 1D, 3D; Table 2)

Lycodontis phasmatodes Smith, 1962: 436 (Type locality: Inhaca Island, Mozambique, western Indian Ocean)

Gymnothorax phasmatodes (Smith, 1962), Böhlke, 1997: 97-98; Hatooka et al., 1998: 2-5; Böhlke et al., 1999: 1649; Böhlke and Smith, 2002: 137.

1. Materials Examined

Six specimens: TOU-AE 0227, 382 mm TL, male, Jul. 2003, Kenting, Pingtung County. TOU-AE 1269, 345 mm TL, female, 08 Aug. 2005; TOU-AE 3271, 334 mm TL, male, 24 Aug. 2006, Chengkung, Taitung County; TOU-AE 3684, 355 mm TL, male, 21 Sept. 2006; TOU-AE 4263, 349 mm TL, male; TOU-AE 4264, 371 mm TL, female, 27 Jul. 2007, Changbin, Taitung County, longline by Captain Jiunn-Shiun Chiou.

2. Diagnosis

An elongate moray with long tapering tail, body yellowish tan, dorsal and anal fins with prominent white margin. Anus

just posterior to midbody. Body elongate with unpatterned coloration. Head long, snout short and blunt, jaws short, eye large. Preanal length 1.8-2.0 (1.9 ± 0.1), tail length 2.0-2.2 (2.1 ± 0.1), trunk length 2.2-2.5 (2.3 ± 0.1), body depth at gill opening 27.8-36.7 (32.4 ± 4.8), body depth at anus 31.6-58.6 (41.8 ± 12.0), predorsal length 13.4-16.1 (14.4 ± 1.2), head length 9.9- 10.4 (10.2 ± 0.3), all in TL. Upper jaw length 2.5-2.8 (2.8 ± 0.2), lower jaw length 2.5-3.1 (2.9 ± 0.3), interorbital width 5.8-9.2 (7.6 ± 1.6), snout length 5.2-5.8 (5.6 ± 0.3), eye diameter 8.9-12.0 (10.3 ± 1.7), all in HL. Premaxillary teeth 6-7 each side, median premaxillary teeth 0-3, maxillary teeth uniserial about 7-12, vomerine teeth uniserial 4-9, dentary teeth uniserial about 12-20 each side (Fig. 3D). Head pores typical, 3 supraorbital pores, 4 infraorbital pores, 6 mandibular pores. Two branchial pores, branchial pores small before gill opening. Gill opening below mid-side. Predorsal vertebrae 5-6, preanal vertebrae 73-77, total vertebrae 165-168; mean vertebral formula 6-75-166.

3. Distribution

Known from the western Indian Ocean and western Pacific, including Mauritius, Mozambique, Moluccas, Japan and Taiwan.

4. Remarks

A moderately small species, the largest recorded was 465 mm TL (the holotype, RUSI 108); we have seen no specimens larger than 382 mm TL (TOU-AE 227) in Taiwan. Sexually mature at 345 mm. Sex was determined for six specimens, two of the study specimens (345 and 371 mm) were females, with eggs of 0.5-0.7 mm, 1.0-1.2 mm in diameter and fecundity 1806 ± 120 eggs; the other four specimens (334-382 mm) were males.

Böhlke (1997) noted that *Gymnothorax phasmatodes* was similar in coloration, dentition and vertebral formula to *G. verrilli* from the eastern Pacific. However, *G. phasmatodes* is different from *G. verrilli* by blunt jaws and paler body color. Hatooka et al. (1998) reported *G. phasmatodes* as a new record from Japan, and described its sexual dimorphism in dentition: the female had three teeth, but the male had no teeth on the mesial part of premaxillary plate. Their result is similar to the counts for our *G. phasmatodes* specimens.

Gymnothorax sagmacephalus Böhlke, 1997
(Figs. 1E, 2B, 3E; Table 2)

Gymnothorax albimarginatus Masuda et al., 1984: plate 28I
(Misidentified)

Gymnothorax sagmacephalus Böhlke, 1997: 100-102. (Type locality: Tokyo Bay, Japan); Böhlke and Smith, 2002: 146.

1. Materials Examined

12 specimens: TOU-AE 0226, 464 mm TL, female, Jul. 2003, Kenting, Pingtung County, longline. TOU-AE 1407, 1409- 1410, 3 specimens, 422-512 mm TL, females, 08 Aug.




Fig. 2. The specimen (TOU-AE 0226, 464 mm TL) of the redescription species, *Gymnothorax sagmacephalus*. A, dorsal view; B, ventral view, showing the coloration pattern.

2005; TOU-AE 2728, 2730, 2732, 3 specimens, 375-482 mm TL, males, 31 Jul. 2006; TOU-AE 2726, 2729, 2731, 3 specimens, 413-426 mm TL. females, 31 Jul. 2006; TOU-AE 5099-5100, 395-468 mm TL, males, 06 July 2008, all were from Taitung County, longline by Captain Jiunn-Shiun Chiou.

2. Diagnosis

An elongate brown moray marked with a dusky saddle on top of head just behind eyes, and a large prominent dark triangle just before the dorsal-fin origin (Fig. 2A). Dorsal and anal fins with a white margin. Body light brownish, pale color on the belly side. Dark brown stripe extends below gill opening along the belly back to anus (Fig. 2B).

3. Description

Preanal length 1.9-2.0 (2.0 ± 0.1), tail length 2.0-2.1 (2.0 ± 0.1), trunk length 2.3-2.5 (2.4 ± 0.1), body depth at gill opening 26.4-34.9 (29.4 ± 2.6), body depth at anus 28.0-45.0 (36.7 ± 4.3), predorsal length 11.4-14.4 (12.9 ± 0.9), head length 8.7-10.3 (9.5 ± 0.5), all in TL. Upper jaw length 2.2-2.9 (2.6 ± 0.2), lower jaw length 2.2-3.1 (2.7 ± 0.2), interorbital width 6.0-8.2 (7.0 ± 0.7), snout length 5.1-6.4 (5.6 ± 0.4), eye diameter 8.5-10.9 (9.8 ± 0.8), all in HL.

Premaxillary teeth 5-7 each side, maxillary teeth uniserial about 6-10, vomerine teeth uniserial 3-8; median premaxillary teeth counts are somewhat different in each sex, 2-3 in female and 0 in male; dentary teeth of mandible in a single row 11-18 on left and 9-17 on right side, it were also different in sex, female 13-18 are more than male 10-14 (Fig. 3E).

Head pores typical, three supraorbital pores, four infraorbital pores, six mandibular pores. Two branchial pores, branchial pores small before gill opening. Gill openings small, just below mid- side of body. Predorsal vertebrae 6-7, preanal vertebrae 74-78, total vertebrae 170-176; mean vertebral formula 6-76-173.

4. Distribution

The species was previously known from Japan, the present record for Taiwan being new.

5. Remarks

A small to moderate-sized moray, the largest known is 534




Fig. 3. Dentitions of the five newly recorded species. A, *Anarchias cantonensis*, TOU-AE 4913 (male, 97 mm TL); B, *Gymnothorax dorsalis*, TOU-AE 4834 (female, 619 mm TL); C, *G. intesi*, TOU-AE 1958 (male, 439 mm TL); D, *G. phasmatodes*, TOU-AE 1269 (female, 189 mm TL); E, *G. sagemcephalus*, TOU-AE 1409 (female, 512 mm TL).

mm TL (the holotype, USNM 149777). Sexes were determined by 12 specimens, eight of the study specimens (395-512 mm TL) were females, four specimens (422-512 mm TL) had ripe eggs with 1.0-1.2 mm egg diameter, and fecundity 3959 ± 622 eggs; the other four specimens (375-482 mm TL) were males.

This is a new record for Taiwan. Hatooka (2002) reported a "*Gymnothorax sagemcephalus* pattern" as a young fish stage of *G. albimarginatus*. This coloration was also illustrated on pl. 28, fig. I in Masuda et al. (1984) and named as *G. albimarginatus* from Japan. It appeared to be *G. sagemcephalus* and showed the prominent triangular saddle on the top of head, as well as the anterior dusky area behind the eyes extending onto the lower jaw. Böhlke (1997) noted that the saddle of *G. sagemcephalus* was so heavily pigmented that it could hardly have been overlooked in any description of the species, and it had not faded in the holotype which was taken 90 years ago. *G. sagemcephalus* could also be differentiated from *G. albimarginatus* by the position of the anus (at mid-body in *sagemcephalus*, well behind mid-body in *albimarginatus*) and the mean vertebral formula (6-76-173 vs. 5-86-187 respectively).

ACKNOWLEDGMENTS

We are very grateful to Captain Jiunn-Shiun Chiou and Ms. Min-Chia Chiang, who helped us to collect these specimens, and two anonymous reviewers for improving the manuscript. These studies were originally supported by National Science Council "NSC 99-2621-B-001-008-MY3; 98-2631-H-019-001"; Fisheries Agency, Council of Agriculture grants "98AS-8.5.2-FA-F1 (2-1)" to Hong-Ming Chen; and University Malaya short research fund "BK027-2011A". Continuing studies of morays have greatly benefitted from discussions with J. E. McCosker (CAS), D. G. Smith (USNM), and J. E. Randall (BPBM). The authors also acknowledge Prof. Chang-Tai Shih for his valuable comments and English editing support.

REFERENCES

- Bloch, M. E. (1795). *Naturgeschichte der ausländischen Fische*. Berlin: Merino and Co. 9, 397-429.
- Böhlke, E. B. (1982). Vertebral formulae of type specimens of eels (Pisces: Anguilliformes). *Proceedings of the Academy of Natural Sciences of Philadelphia* 134, 31-49.
- Böhlke, E. B. (1997). Notes on the identity of elongate unpatterned Indo-Pacific morays, with description of a new species (Muraenidae, Subfamily Muraeninae). *Proceedings of the Academy of Natural Sciences of Philadelphia* 147, 89-109.
- Böhlke, E. B. and J. E. McCosker (2001). The moray eels of Australia and New Zealand, with the description of two new species (Anguilliformes: Muraenidae). *Records of the Australian Museum* 53(1), 71-102.
- Böhlke, E. B. and J. E. Randall (2000). A review of the moray eels (Anguilliformes: Muraenidae) of the Hawaiian Island, with descriptions of two new species. *Proceedings of the Academy of Natural Sciences of Philadelphia* 150, 203-278.
- Böhlke, E. B. and D. G. Smith (2002). Type catalogue of Indo-Pacific Muraenidae. *Proceedings of the Academy of Natural Sciences of Philadelphia* 152, 89-172.
- Böhlke, E. B., J. E. McCosker, and D. G. Smith (1999). Family Muraenidae." In Carpenter K.E. and Niem V.N. (eds.), *FAO species identification guide for fishery purposes. The living marine resources of the Western Central Pacific, Batoid fishes, chimaeras, bony fishes* 6(1), 1643-1657.
- Chen, H. M. and K. H. Loh (2007). *Gymnothorax shaoi*, a new species of moray eel (Anguilliformes: Muraenidae) from southeastern Taiwan. *Journal of Marine Science and Technology* 15, 76-81.
- Chen, H. M., K. H. Loh, and K. T. Shao (2008). A new species of moray eel, *Gymnothorax taiwanensis* (Anguilliformes: Muraenidae) from eastern Taiwan. *The Raffles Bulletin of Zoology, Supplement* 19, 131-134.
- Chen, H. M., K. T. Shao, and C. T. Chen (1994). A review of the muraenid eels (Family Muraenidae) from Taiwan with descriptions of twelve new records. *Zoological studies* 33(1), 44-64.
- Chen, H. M., K. T. Shao, and C. T. Chen (1997). Systematic Studies on the Muraenid Fishes (Muraenidae) from the Waters Around Taiwan. Ph. D. Thesis, Department of Fishery Science, National Taiwan Ocean University, Keelung, 1-257. (In Chinese)
- Fourmanoir, P. and I. Rivaton (1979). Poissons de la pente récifale externe de Nouvelle-Calédonie et des Nouvelles-Hébrides. *Cahiers de l'Indo-Pacifique* 1(4), 405-443.
- Hatooka, K. (1986). Sexual dimorphism found in teeth of three species of moray eels. *Japanese Journal of Ichthyology* 32(4), 379-386.
- Hatooka, K. (2002). Family Muraenidae. In Nakabo, T., (ed), *Fishes of Japan with pictorial keys to the species*. English edition. Tokai University Press.
- Hatooka, K., W. Hiramatsu, and H. Ikeda (1998). First record of a muraenid eel *Gymnothorax phasmatodes* (Smith, 1962) from Japan. *I. O. P. Diving News* 9(5), 2-5.

- Jordan, D. S. and A. Seale (1906). The fishes of Samoa. Description of the species found in the archipelago, with a provisional check-list of the fishes of Oceania. Bulletin of the Bureau Fisheries 25, 173-455.
- Linnaeus, C. (1758). Systema Naturae. Ed. X. (Systema naturae per regna tria naturae, secundum classes, ordines, genera, species, cum characteribus, differentiis, synonymis, locis. Tomus I. Editio decima, reformata.) Holmiae. 1, 1-824.
- Loh, K. H., I. S. Chen, J. E. Randall, and H. M. Chen (2008). A review and molecular phylogeny of the moray eel subfamily Uropterygiinae (Anguilliformes: Muraenidae) from Taiwan, with description of a new species. The Raffles Bulletin of Zoology, Supplement 19, 135-150.
- Loh, K. H., K.T. Shao, and H. M. Chen (2011). *Gymnothorax melanosomatus*, a new moray eel (Teleostei: Anguilliformes: Muraenidae) from south-eastern Taiwan. Zootaxa 3134, 43-52.
- Loh, K. H., K.T. Shao, and H. M. Chen (2012). Additions to the Taiwan marine eel fauna with first records of three rare moray eels (Anguilliformes: Muraenidae). Journal of marine science and technology 20(2), 210-215.
- Masuda, H., K. Amaoka, C. Araga, T. Uyeno, and T. Yoshino (1984). The Fishes of The Japanese Archipelago. Tokai Univ. Press. 1-437.
- McCosker, J. E. and A. L. Stewart (2006). Additions to the New Zealand marine eel fauna with the description of a new moray, *Anarchias supremus* (Teleostei: Muraenidae), and comments on the identity of *Gymnothorax griffini* Whitley and Phillips. Journal of the Royal Society of New Zealand 36(2), 83-95.
- Reece, J. S., D. G. Smith, and E. Holm (2010). The moray eels of the *Anarchias cantonensis* group (Anguilliformes: Muraenidae), with description of two new species. Copeia 3, 421-430.
- Rüppell, W. P. E. S. (1838). Neue Wirbelthiere zu der Fauna von Abyssinien gehörig. Fische des Rothen Meeres, Frankfurt-am-Main, 148.
- Schultz, L. P. (1943). Fishes of the Phoenix and Samoan Islands collected in 1939 during the expedition of the U.S.S. "Bushnell". U.S. National Museum Bulletin 180, 1-316.
- Seale, A. (1917). New species of apodal fishes. Bulletin of the Museum of Comparative Zoology 61(4), 79-94.
- Shao, K. T., H. C. Ho, P. L. Lin, P. F. Lee, M. Y. Lee, C. Y. Tsai, Y. C. Liao, Y. C. Lin, J. P. Chen, and H. M. Yeh (2008). A checklist of the fishes of southern Taiwan, northern South China Sea. The Raffles Bulletin of Zoology, Supplement 19, 233-271.
- Smith, D. G. (1994). Catalog of type specimens of recent fishes in the National Museum of Natural History, Smithsonian Institution, 6: Anguilliformes, Saccopharyngiformes, and Notacanthiformes (Teleostei: Elopomorpha). Smithsonian Contributions to Zoology 566, 50.
- Smith, J. L. B. (1962). The moray eels of the Red Sea and western Indian Ocean. Ichthyological Bulletin J. L. B. Smith Institute of Ichthyology 23, 421-444.

ELECTRONIC REFERENCE

- Eschmeyer, W. N. and J. D. Fong (2013). Species by family/subfamily. Catalog of Fishes, California Academy of Sciences," On-line version dated 25 March 2013. <http://research.calacademy.org/redirect?url=http://research-archive.calacademy.org/research/ichthyology/catalog/index.asp>