



GYMNOTHORAX SHAOI, A NEW SPECIES OF MORAY EEL (ANGUILLIFORMES: MURAENIDAE) FROM SOUTHEASTERN TAIWAN

Hong-Ming Chen

*Department of Aquaculture, National Taiwan Ocean University, Keelung, Taiwan 202, R.O.C.,
hmchen@mail.ntou.edu.tw*

Kar-Hoe Loh

Department of Aquaculture, National Taiwan Ocean University, Keelung, Taiwan 202, R.O.C.

Follow this and additional works at: <https://jmstt.ntou.edu.tw/journal>



Part of the [Aquaculture and Fisheries Commons](#)

Recommended Citation

Chen, Hong-Ming and Loh, Kar-Hoe (2007) "GYMNOTHORAX SHAOI, A NEW SPECIES OF MORAY EEL (ANGUILLIFORMES: MURAENIDAE) FROM SOUTHEASTERN TAIWAN," *Journal of Marine Science and Technology*.

Vol. 15: Iss. 2, Article 2.

DOI: 10.51400/2709-6998.2035

Available at: <https://jmstt.ntou.edu.tw/journal/vol15/iss2/2>

This Research Article is brought to you for free and open access by Journal of Marine Science and Technology. It has been accepted for inclusion in Journal of Marine Science and Technology by an authorized editor of Journal of Marine Science and Technology.

GYMNOTHORAX SHAOI, A NEW SPECIES OF MORAY EEL (ANGUILLIFORMES: MURAENIDAE) FROM SOUTHEASTERN TAIWAN

Acknowledgements

We wish to thank Dr. Kwang-Tsao Shao, Acting Director of the Research Center for Biodiversity, Academia Sinica; and Dr. Ching-Fong Chang in our University, for their valuable advice and support throughout the studies. We give particular thanks to Dr. Daniel J. Sheehy and Dr. Susan Vik. for revising the manuscript. We are very grateful to Mr. Wung-Jyh Yang and Captain Jiunn-Shiun Chiou who collected the type and relative specimens. Mr. Wei-Jen Chen, Mr. Yen-Chieh Tseng, Mr. Chao-Sen Yang and Miss Shih-Ying Tien measured the type specimens and reared the comparative morays in our aquaria. These studies were originally supported by Fisheries Agency, Council of Agriculture grants "92AS-9.1.1-FA-F1(8); 93AS-9.1.1-FA-F2(1-4); 94AS-9.4.1-FA-F1 (3-4)" to Hong-Ming Chen. This study was also partially supported by the Center for Marine Bioscience and Biotechnology in National Taiwan Ocean University.

GYMNOTHORAX SHAOI, A NEW SPECIES OF MORAY EEL (ANGUILLIFORMES: MURAENIDAE) FROM SOUTHEASTERN TAIWAN

Hong-Ming Chen* and Kar-Hoe Loh*

Key words: new species, Muraenidae, moray eel, *Gymnothorax shaoi*.

ABSTRACT

A new species of moray eel, *Gymnothorax shaoi*, is described here based on ten specimens collected in southeastern Taiwan. The new muraenid is distinct from its most similar species, *Gymnothorax fimbriatus*, in the character combinations of having brown body color, dark brown spots in at least three rows on body side and spots on posterior dorsal and caudal fin not forming oblique streaks; paler rictus, jaws, and inner mouth; fins without pale margins; no bend in lower jaw; fewer total vertebrae; and smaller in maximum body size. *G. shaoi*'s eyes have yellow to orange irises, body with dark reddish brown spots, and no yellow mucus on the top of head when alive. *G. shaoi* also differs from another similar species, *G. reevesii*, in several characteristics: smaller body depth, longer snout length, absence of pale posterior nostrils, with a few dark brown dots on the dorsal fin, no blackish margins on fins, and has more inner maxillary and inner dentary teeth.

INTRODUCTION

During 2001-2003 in a series of projects entitled "Investigations and studies of endangered muraenid fishes from the waters around Taiwan", we collected many specimens of moray eels from the waters surrounding Taiwan. Among these, we noted some exceptional specimens that had a color pattern similar to but distinct from *Gymnothorax fimbriatus* (Bennett, 1832) [1]. In June 2003, a live specimen with variable patterned "*G. fimbriatus*" was obtained from Chengkung in southeastern Taiwan. It was transported to our laboratory in the Department of Aquaculture at National Taiwan Ocean University, and maintained alive to observe its behavior and morphological characters in detail. We found that this specimen was quite different from *G. fimbriatus* and could be an undescribed species. In August 2003, a fishing shop

master, Mr. Wung-Jyh Yang, provided 8 additional specimens of this *Gymnothorax* sp. from Chengkung. In August 2005, Captain Jiunn-Shiun Chiou provided another live specimen of this species from Changbin in southeastern Taiwan. It was kept alive and recorded in detail as well in our laboratory for more than 4 months. After examination of these specimens and make the comparison on their morphology to other similar moray species, we propose and describe it here as a new species.

MATERIALS AND METHODS

The methods of measurement followed those of Chen *et al.* [8] or Böhlke and Randall [5]. Proportional measurements of type specimens of the new moray were expressed as percentage of the total length (TL) or the head length (HL). Preanal length (PAL) was measured from the snout tip to the mid-anus; body depth was measured at the gill openings (DGO) and at the anus (DA) that did not include the fins; snout length was measured from snout tip to the anterior margin of the eye; upper jaw length was from snout tip to mouth angle; and lower jaw length was from lower jaw tip to mouth angle. Counts for the vertebral formula were obtained from radiographs, as explained in Böhlke [3] and Chen *et al.* [9]; the mean vertebral formula (MVF) provided mean values for predorsal-preanal-total vertebrae counts. Teeth counts referenced Hatooka [11], were approximate, and included sockets of missing teeth. Gonadal type was determined by gross and histological examination of the muraenid gonad.

The type specimens were stored in the Museum of the Research Center for Biodiversity, Academia Sinica (ASIZP), and in the collections of the Laboratory of Aquatic Ecology, Department of Aquaculture, National Taiwan Ocean University (TOU-AE). The related muraenid specimens that we examined were also from National Museum of Marine Science & Technology (NMMSTP) and Fisheries Research Institute (FRI).

Paper Submitted 03/27/06, Accepted 09/15/06. Author for Correspondence: Hong-Ming Chen. E-mail: hmchen@mail.ntou.edu.tw.

*Department of Aquaculture, National Taiwan Ocean University, Keelung, Taiwan 202, R.O.C.

RESULTS AND DISCUSSION

Gymnothorax shaoi n. sp.

(Chinese name: 邵氏裸胸鯨; Shao's Rou-hsun-tsung)

(English name: Shao's moray eel)

(Figures 1, 2, 3 a, 4; Table 1)

Holotype: ASIZP0062978 (Figure 1), male, 567 mm TL, off shore of Sanshengtai (23.130°N, 121.414°E), Chengkung, Taitung, longline, Wung-Jyh Yang, 19 Aug. 2003.

Paratypes: ASIZP0062979-80, 2 specimens, females, 401-457 mm TL, off shore of Sanshengtai, longline, Wung-Jyh Yang, 19 Aug. 2003; TOU-AE0370, 0371, 0373, 0375, 4 specimens, males, 398-608 mm TL, and TOU-AE0372, female, 405 mm TL, off shore of Sanshengtai, longline, Wung-Jyh Yang, 19 Aug. 2003; TOU-AE0381 (Figure 3a), female, 379 mm TL, tidal pool of Hsiaogang (23.156°N, 121.405°E), Chengkung, Taitung, clove oil, Kar-Hoe Loh, 22 Jun. 2003; TOU-AE1797 (Figure 4), female, 492 mm TL, off shore of Changbin (23.311°N, 121.453°E), Taitung, longline, Captain Jiunn-Shiun Chiou, 1 Aug. 2005.

Comparative materials:

***Gymnothorax fimbriatus*:** 77 specimens. ASIZP0056652, 2: 171-195 mm; NMMSTP00216, 121 mm; FRIP20092, 610 mm; TOU-AE0211, 764 mm; TOU-AE0218, 633 mm; TOU-AE0288, 488 mm; TOU-AE0289, 423 mm; TOU-AE0391, 337 mm; TOU-AE0392, 305 mm; TOU-AE0504, 113 mm; TOU-AE0505, 246 mm; TOU-AE0506, 285 mm; TOU-AE0530, 599 mm; TOU-AE0569, 340 mm; TOU-

AE0631-0637, 7: 332-842 mm; TOU-AE0650-0651, 2: 224-351 mm; TOU-AE0654-0656, 3: 246-694 mm; TOU-AE0727-0734, 8: 351-770 mm; TOU-AE0802, 497 mm; TOU-AE0808, 395 mm; TOU-AE0811-0813, 3: 333-652 mm; TOU-AE0921, 416 mm; TOU-AE0931, 405 mm; TOU-AE0936-0937, 2: 427-605 mm; TOU-AE0939-0940, 2: 408-599 mm; TOU-AE0943, 711 mm; TOU-AE1160, 806 mm; TOU-AE1518-1521, 4: 510-664 mm; TOU-AE1527-1531, 5: 413-652 mm; TOU-AE1534-1538, 5: 334-689 mm; TOU-AE1544-1549, 6: 383-883 mm; TOU-AE1665-1669, 5: 457-669 mm; TOU-AE1674-1677, 4: 348-631 mm.

***Gymnothorax reevesii*:** 9 specimens. ASIZP0056653, 650 mm; NMMSTP00941, 698 mm; TOU-AE0130, 432 mm; TOU-AE0219, 534 mm; TOU-AE0581, 685 mm; TOU-AE0831-0833, 3: 457-592 mm; TOU-AE1369, 725 mm.

Diagnosis:

Body length moderate (maximum size reaches 608 mm TL), smaller body depth (Figure 1). Dorsal fin moderately high; origin before the gill opening. Anus close to mid-body point. Jaw teeth caniniform and generally uni-serial, but bi-serial on anterior dentaries and maxillae. Body and fins light brown, with larger dark brown spots on body side in three rows at minimum, with only a few dark brown dots on the dorsal fin; paler jaws, rictus, and abdomen; anterior area of head lacks large dark spots. When alive, the irises of the eyes are yellow to orange and the body spots are dark reddish brown.



Fig. 1. Holotype of *Gymnothorax shaoi* n. sp., male, ASIZP 0062978, 567 mm TL, Chengkung, Taitung, longline.




Fig. 2. Dentitions of type specimens of *Gymnothorax shaoi*: holotype, male; (a) ASIZP 0062978; paratypes, females; (b) ASIZP 0062980; (c) TOU-AE 0381.

Description:

Data for holotype, followed in parentheses by the mean and range of data for the holotype and eight paratypes (exclude specimen TOU-AE 1797). The asterisk (*) indicates that data were affected by a missing tail-tip. Tail length 2.02 (2.00*; 1.94*-2.06); trunk length 2.70 (2.67; 2.56-2.81); depth at anus 17.26 (20.71*; 17.26-24.62); depth at gill opening 14.32 (17.39*; 14.32-23.53); head length 7.46 (7.77*; 7.35*-8.26), all in TL. Predorsal length 1.58 (1.54; 1.45-1.76); upper jaw length 2.54 (2.38; 2.19-2.54); lower jaw length 2.55 (2.40; 2.24-2.55); snout length 5.91 (5.70; 5.04-6.19); eye diameter 13.06 (11.71; 10.25-13.06); interorbital width 7.68 (7.72; 5.98-9.72), all in HL. Predorsal vertebrae 5 (3.9; 3-5); pre-epipleural vertebrae 52 (49.9; 45-53); preanal vertebrae 57 (55.8; 53-57); and total vertebrae 129 (125.1*; 112*-130). Lengths, proportions, vertebral counts and gonadal type of the holotype and eight paratypes of *G. shaoi* n. sp. are shown in Table 1.

Dorsal fin moderately high, origin before the gill opening and above the 3-5th vertebra. The depth of anal fin low, origin just behind anus and below 53-57th vertebra. Gill opening on mid-body, and its length nearly equal to eye diameter.

Anterior nostrils were tubular on each side of snout tip. Tips of anterior nostrils were lower and with darkish flounced rims. Posterior nostrils were short tubular over each front edge of eye with darkish raised funneled rims.

Head pores distinct. Supraorbital canal with three pores, one of them situated anteroventrally to anterior nostril. Infraorbital canal with four pores. Mandibular canal with six to seven pores. Two branchial pores situated near the mid-point area of origin of dorsal fin to gill opening.

Mouth in terminal position and closing completely. Caniniform teeth in jaws, pointed and slightly retrorse



(a)



(b)



(c)

Fig. 3. Comparisons of head characters of *Gymnothorax shaoi* (a: paratype, TOU-AE 0381) with two similar species, *G. fimbriatus* (b) and *G. reevesii* (c).



Fig. 4. The paratype (TOU-AE1797, 492 mm TL, from Changbin) of *Gymnothorax shaoi* reared in our laboratory exhibited the paler ground color at night.

(Figure 2). Teeth of peripheral series of premaxillary plate 14-23 (these counts include shed teeth); several small teeth found in some spaces between lateral teeth of peripheral series of premaxillary plate. Mesial series of premaxillary plate with 3-4 teeth, with posterior larger. Vomerine teeth small, in a straight series and 5-9 in number. Maxillary teeth in double rows, 17-18 canines on outer row of each side and 3-4 longer inner canines in anterior maxilla. Mandibular teeth in double rows, 4-5 larger canines on inner row of anterior dentary and 19-23 canines on outer row of dentary. As in the peripheral series of premaxillary teeth, several small teeth found in the space between larger teeth on anterior dentary near symphysis.

Body color in formalin or alcohol grey brown. Dark brown spots on body side mainly in three rows, one row along dorsal fin base, one row along lateral-line, and one row along anal fin base. Some small dark brown dots were scattered on dorsal fin or body side. Paler rictus and abdomen. No obviously dark spots on jaws, inner mouth, chin and anal fin. Body color when alive similar to color in preservative, but the brown ground color a little reddish at day (Figure 3a) or more pale at night (Figure 4). When alive, *G. shaoi* has eyes with yellow to orange irises and body with dark reddish brown spots.

Distribution:

From southeastern Taiwan. Off shore of Chengkung (23.130°N, 121.414°E) to Changbin (23.311°N, 121.453°E), Taitung County, Taiwan.

Remarks:

The new muraenid *Gymnothorax shaoi* is clearly distinct from the most similar species *G. fimbriatus* (Bennett) [1] (Figure 3b) in the character combinations of brown body color, dark brown spots on body side in at least three rows, spots on posterior dorsal and caudal fin that do not form oblique streaks; paler rictus, jaws and inner mouth; fins without pale margins; no bend in lower jaw; fewer total vertebrae (mean total vertebrae 128 vs. 143 [6]); smaller maximum size (reaches 608 vs. 883 mm TL) (Table 2). *G. shaoi* has eyes with yellow to orange irises, body with dark reddish brown spots, and lack yellow mucus character on the top of head when alive. *G. fimbriatus* has whitish to yellowish body color and dense yellow mucus on head, blackish spots on body, blackish mouth corners, dark jaws, blackish skin in inner mouth, remarkable whitish margin on the anal and caudal fins, and dark reddish eye irises when alive.

G. shaoi also differs from another similar species *G. reevesii* (Richardson) [12] (Figure 3c) in the characters of smaller body depth (DGO. 4.3-7.0 vs 9.1-10.1 % of TL.) [7, 9], longer snout length (16.2-19.9 vs 13.5-16.1 % of HL.), absence of pale posterior nostrils, only a few dark brown dots on the dorsal fin, and absence of blackish margins on fins. *G. reevesii* has numerous large obscure dark brownish spots in 3-5 rows along the body side, many small brownish dots speckled about the body, posterior caudal fin with oblique comb-like dark brown bands, fins with a dark margin. *G. shaoi* has more inner maxillary (3-4 vs. 0) and inner dentary (4-5

Table 1. Lengths, proportions, vertebral counts and gonadal type of the holotype and eight paratypes of *Gymnothorax shaoi* n. sp.

	Holotype		Paratypes									
	ASIZP0062978 (mm) (%)	ASIZP0062979 (mm) (%)	ASIZP0062980 (mm) (%)	TOU-AE0370 (mm) (%)	TOU-AE0371 (mm) (%)	TOU-AE0372 (mm) (%)	TOU-AE0373 (mm) (%)	TOU-AE0375 (mm) (%)	TOU-AE0381 (mm) (%)			
Total length	567	457	401	458	416	405*	398*	608	379			
% of total length												
Tail length	281 49.6	236 51.6	203 50.6	224 48.9	213 51.2	201 49.6*	193 48.5*	303 49.8	191 50.4			
Trunk length	210 37.0	169 37.0	148 36.9	179 39.1	157 37.7	153 37.8*	152 38.2*	233 38.3	135 35.6			
Body depth at anus	32.9 5.8	22.0 4.8	16.3 4.1	23.2 5.1	23.0 5.5	22.3 5.5*	20.2 5.1*	25.4 4.2	15.7 4.1			
Body depth at gill opening	39.6 7.0	28.0 6.1	23.5 5.9	26.0 5.7	24.3 5.8	23.0 5.7*	23.8 6.0*	37.2 6.1	16.1 4.3			
Head length	76.0 13.4	58.0 12.7	52.2 13.0	55.4 12.1	51.6 12.4	55.1 13.6*	51.4 12.9*	78.0 12.8	49.3 13.0			
% of head length												
Predorsal length	48.3 63.5	38.1 65.7	36.0 69.0	37.4 67.5	35.4 68.6	31.3 56.8	34.6 67.4	49.7 63.7	32.0 64.9			
Upper Jaw length	29.9 39.3	26.4 45.6	20.6 39.4	24.1 43.4	23.6 45.7	23.1 41.8	21.0 40.8	31.5 40.4	21.0 42.6			
Lower Jaw length	29.8 39.3	25.9 44.7	21.2 40.7	23.5 42.4	21.7 42.1	24.2 43.9	20.3 39.5	32.0 41.0	21.1 42.8			
Snout length	12.9 16.9	11.5 19.8	8.4 16.2	9.3 16.8	9.8 19.0	9.6 17.3	8.7 16.9	13.6 17.5	8.9 18.0			
Eye diameter	5.8 7.7	5.2 9.0	5.1 9.8	4.7 8.5	4.7 9.1	4.3 7.9	4.7 9.1	6.1 7.8	4.2 8.5			
Interorbital width	9.9 13.0	9.7 16.7	6.5 12.4	5.7 10.3	7.4 14.4	7.8 14.1	5.5 10.7	10.0 12.8	7.2 14.6			
Predorsal vertebrae	5	4	4	4	4	3	4	3	4			
Preanal vertebrae	57	56	56	57	53	53	56	57	57			
Total vertebrae	129	130	128	128	125	112*	118*	129	127			
Gonadal type	testis	ovary	ovary	testis	testis	ovary	testis	testis	ovary			

Note: *Tip of tail missing

Table 2. Comparisons of proportions, counts, coloration, and other characters among *Gymnothorax fimbriatus*, *G. reevesii* and *G. shaoi* n. sp.

	<i>G. fimbriatus</i>	<i>G. reevesii</i>	<i>G. shaoi</i>
% of total length			
Body depth at gill opening	5.9-8.1	9.1-10.1	4.3-7.0
% of head length			
Snout length	14.9-18.9	13.5-16.1	16.2-19.9
Upper Jaw length	34.5-57.1	38.7-45.1	39.3-45.8
Lower Jaw length	33.8-58.1	37.1-43.4	39.3-44.7
Counts			
Total vertebrae (mean)	131-133(132); 143*	125-128 (127)	125-130 (128)
Inner maxillary teeth	1-4	0	3-4
Inner dentary teeth	2-4	0	4-5
Coloration			
Body color	Whitish to yellowish	Yellowish brown to brownish	Gray to brownish
Body spots	Blackish	Dark brownish	Reddish brown
Irises of eyes	Dark reddish	Dark reddish	Yellowish to orangey
Jaws	Darkish	Brownish	Pale
Margins of fins	Whitish	Dark brownish	Brownish
Posterior nostrils	Brownish	Pale	Darkish
Rictus of mouth	Blackish	Brownish	Pale
Skin of inner mouth	Blackish	Brownish	Pale
Top of head yellowish	Yes	No	No
Bend in lower jaw	Yes	No	No
Maximum total length	883 mm	698 mm	608 mm

Note: *Data from [6]

vs. 0) teeth than *G. reevesii*.

Böhlke and Smith [6] reviewed the type catalogue of Indo-Pacific Muraenidae. They remarked on Bleeker's specimen (BMNH 1867.11.28.238) of "*Gymnothorax bullatus*" [2] (misidentified, not *Muraena bullata* Richardson [13]) as the holotype of "*Muraena microspila* Günther[10]". Color patterns of the specimen depicted in Bleeker's Atlas [2] was somewhat similar to our *G. shaoi*. However, the specimen had darkish fin margins and its dentition was closer to the diagnosis of *G. reevesii* which was reviewed in Chen *et al.* [7]. Böhlke *et al.* [4] cited the drawing of "*Gymnothorax bullatus*" from Bleeker Atlas [2] and renamed it as *G. reevesii*. Böhlke and Smith [6] also remarked that the specimen (BMNH 1867.11.28.238) was *G. reevesii*, and considered "*Muraena microspila*" as the junior synonym of *G. reevesii*.

Key to *Gymnothorax* species which have numerous dark spots speckling on gray to brownish body:

- 1a. Margin of fins yellowish *G. buroensis*
 1b. Margin of fins not yellowish 2

- 2a. Body with black round spots; spots are more dense and numerous on head *G. favagineus*
 2b. Body with dark spots; spots are not dense and numerous on head 3
 3a. Upper region of head yellowish *G. fimbriatus*
 3b. Upper region of head not yellowish 4
 4a. Dark reddish irises, caudal fin with darkish margin *G. reevesii*
 4b. Yellowish to orangey irises, caudal fin without darkish margin *G. shaoi*

Etymology:

We name this species *Gymnothorax shaoi* in honor of Dr. Kwang-Tsao Shao, who has contributed greatly to establish the Taiwanese fish database as well as supervise and support for our muraenid studies.

ACKNOWLEDGEMENTS

We wish to thank Dr. Kwang-Tsao Shao, Acting Director of the Research Center for Biodiversity,

Academia Sinica; and Dr. Ching-Fong Chang in our University, for their valuable advice and support throughout the studies. We give particular thanks to Dr. Daniel J. Sheehy and Dr. Susan Vik. for revising the manuscript. We are very grateful to Mr. Wung-Jyh Yang and Captain Jiunn-Shiun Chiou who collected the type and relative specimens. Mr. Wei-Jen Chen, Mr. Yen-Chieh Tseng, Mr. Chao-Sen Yang and Miss Shih-Ying Tien measured the type specimens and reared the comparative morays in our aquaria.

These studies were originally supported by Fisheries Agency, Council of Agriculture grants "92AS-9.1.1-FA-F1(8); 93AS-9.1.1-FA-F2(1-4); 94AS-9.4.1-FA-F1(3-4)" to Hong-Ming Chen. This study was also partially supported by the Center for Marine Bioscience and Biotechnology in National Taiwan Ocean University.

REFERENCES

1. Bennett, E.T., "Observations on a Collection of Fishes from the Mauritius, Presented by Mr. Telfair, with Characters of New Genera and Species," *Proceedings of the Zoological Society of London*, Vol. 1, pp. 165-169 (1832).
2. Bleeker, P., "Murènes, Synbranches, Leptocéphales," *Atlas Ichthyologique des Indes Orientales Néerlandaises, Publie Sous les Auspices du Gouvernement Colonial Néerlandaises. Tome IV*, Amsterdam, pp. 1-150, pls. 144-193 (1864).
3. Böhlke, E.B., "Vertebral Formulae of Type Specimens of Eels (Pisces: Anguilliformes)," *Proceedings of the Academy of Natural Sciences of Philadelphia*, Vol. 134, pp. 31-49 (1982).
4. Böhlke, E.B., McCosker, J.E., and Smith, D.G., "Muraenidae-Morays," in K.E. Carpenter and V.H. Niem (eds.), *FAO Species Identification Guide for Fishery Purposes. The Living Marine Resources of the Western Central Pacific*, Rome: FAO of the United Nations, Vol. 3, pp. 1643-1657 (1999).
5. Böhlke, E.B. and Randall, J.E., "A Review of the Moray Eels (Anguilliformes: Muraenidae) of the Hawaiian Island, with Descriptions of Two New Species," *Proceedings of the Academy of Natural Sciences of Philadelphia*, Vol. 150, pp. 203-278 (2000).
6. Böhlke, E.B. and Smith, D.G., "Type Catalogue of Indo-Pacific Muraenidae," *Proceedings of the Academy of Natural Sciences of Philadelphia*, Vol. 152, pp. 89-172 (2002).
7. Chen, H.M., Shao, K.T., and Chen, C.T., "A Review of the Muraenid Eels (Family Muraenidae) from Taiwan with Descriptions of Twelve New Records," *Zoological Studies*, Vol. 33, No. 1, pp. 44-64 (1994).
8. Chen, H.M., Shao, K.T., and Chen, C.T., "A New Moray Eel, *Gymnothorax niphostigmus*, (Anguilliformes: Muraenidae) from Northern and Eastern Taiwan," *Zoological Studies*, Vol. 35, No. 1, pp. 20-24 (1996).
9. Chen, H.M., Chen, C.T., and Shao, K.T., "Systematic Studies on the Muraenid Fishes (Muraenidae) from the Waters Around Taiwan," Ph. D. Thesis, Department of Fishery Science, National Taiwan Ocean University, Keelung, Taiwan, pp. 1-257 (1997) (in Chinese).
10. Günther, A., *Catalogue of the Fishes in the British Museum*, British Museum, London, Vol. 8, pp. 1-549 (1870).
11. Hatooka, K., "Sexual Dimorphism Found in Teeth of Three Species of Moray Eels," *Japanese Journal of Ichthyology*, Vol. 32, No. 4, pp. 379-386 (1986).
12. Richardson, J., "Ichthyology," In Hinds, R.B. (eds.), *The Zoology of the Voyage of H.M.S. Sulphur, Under the Command of Captain Sir Edward Belcher, etc., During the Years 1836-42*, Smith, Elder and Co., London, pp. 51-150, pls. 35-64 (1844-1845).
13. Richardson, J., *Ichthyology of the Voyage of H.M.S. Erebus and Terror, under the Command of Captain Sir JC Ross, etc., During the Years 1839-43*, London, Vol. 2, pp. 1-139 (1844-1848).