



Journal of Marine Science and Technology



Volume 30
Issue 5 Review-Special Issue (Part 2)

Article 23

Key to the species of Augaptilidae Haloptilus occurring in the China Seas

Chang-tai Shih

National Taiwan Ocean University, ctshih@mail.ntou.edu.tw

Qing-Chao Chen

South China Sea Institute of Oceanology

Yang-Chi Lan

Fisheries Research Institute

Shih-Hui Hsiao

National Taipei University of Education

Chi-Yu Weng

Institute of Oceanography, National Taiwan University/Ocean Data Bank of the Ministry of National Science and Technology Council, Taiwan

Follow this and additional works at: <https://jmstt.ntou.edu.tw/journal>



Part of the Fresh Water Studies Commons, Marine Biology Commons, Ocean Engineering Commons, Oceanography Commons, and the Other Oceanography and Atmospheric Sciences and Meteorology Commons

Recommended Citation

Shih, Chang-tai; Chen, Qing-Chao; Lan, Yang-Chi; Hsiao, Shih-Hui; and Weng, Chi-Yu (2022) "Key to the species of Augaptilidae Haloptilus occurring in the China Seas," *Journal of Marine Science and Technology*: Vol. 30: Iss. 5, Article 23.

DOI: 10.51400/2709-6998.2614

Available at: <https://jmstt.ntou.edu.tw/journal/vol30/iss5/23>

This Review-Taxonomic Index is brought to you for free and open access by Journal of Marine Science and Technology. It has been accepted for inclusion in Journal of Marine Science and Technology by an authorized editor of Journal of Marine Science and Technology.

REVIEW-TAXONOMIC INDEX

Key to the Species of Augaptilidae *Haloptilus* Occurring in the China Seas

Chang-tai Shih ^{a,*}, Qing-Chao Chen ^b, Yang-Chi Lan ^c, Shih-Hui Hsiao ^d, Chi-Yu Weng ^e

^a National Taiwan Ocean University, Taiwan

^b South China Sea Institute of Oceanology, China

^c Fisheries Research Institute, Taiwan

^d National Taipei University of Education, Taiwan

^e Institute of Oceanography, National Taiwan University/Ocean Data Bank of the Ministry of National Science and Technology Council, Taiwan

Haloptilus acutifrons(4a/21a/f), *austini*(12a/f) (male unknown), *chierchiai*(7a/18a/f), *fertilis*(22a/f) (female unknown), *fons*(7b/f) (male unknown), *longicirrus*(14a/f) (male unknown), *longicornis*(13a/16a/f), *mucronatus*(5a/15a/f), *ornatus*(9a/22b/f), *oxycephalus*(3a/22b/f), *paralongicirrus*(14b/18b/f), *plumosus*(11a/f) (male unknown), *spinosiceps*(5b/19a/f), *tenuis*(11b/22c/f) (**Haloptilus bulliceps* Is excluded from this key because only copepodid IV has been reported and illustrated (Farran, 1926; Rose, 1933; Saraladevi, 1977; Lian & Qian, 1984; Zhang, Zhao et al., 2010; Lian et al., 2018)).

1a	Female	2
1b	Male	15
2a/1a	Front of cephalosome with pointed spine	3
2b	Front of cephalosome rounded.....	6
3a/2a	Frontal spine of cephalosome long and straight.....	<i>Haloptilus oxycephalus</i>
3b	Frontal spine of cephalosome short and curved downward in lateral view.....	4
4a/3b	Antennule, when pressed against body, exceeding distal margin of caudal ramus by last 4 segments. Maxillule, endopod with 4 setae. Prosome/uromosome length ratio less than 6.....	<i>Haloptilus acutifrons</i>
4b	Antennule, when pressed against body, exceeding distal margin of caudal ramus by last 2 segments. Maxillule, endopod 1-segmented, with 3 setae. Prosome/uromosome length ratio larger than 7	5
5a/4b	Maxillule, exopod with 9 setae. Prosome length/width ratio larger than 4.5. Prosome/uromosome length ratio larger than 8.5	<i>Haloptilus mucronatus</i>
5b	Maxillule, exopod with 10 setae. Prosome length/width ratio smaller than 3.5. Prosome/uromosome length ratio smaller than 7.5.....	<i>Haloptilus spiniceps</i>
6a/2b	Maxillule, endopod 2-segmented	7
6b	Maxillule, endopod 1-segmented	8
7a/6a	Antenna endopod/exopod length ratio 2.7. Maxillule, endopod segment 1 with 3, segment 2 with 4 setae; praecoxal arthrite with 7 setae. Prosome length/width ratio larger than 3	<i>Haloptilus chierchiai</i>
7b	Antenna endopod/exopod length ratio 1.5. Maxillule, endopod segment 1 with 4, segment 2 with 8 setae; praecoxal arthrite with 10–11 setae. Prosome length/width ratio smaller than 3	<i>Haloptilus fons</i>
8a/6b	Maxillule: endopod with 1 seta.....	<i>Haloptilus belliceps</i>
8b	Maxillule: endopod with more than 1 seta	9
9a/8b	Maxillule, endopod 1-segmented, with 2 setae	<i>Haloptilus ornatus</i>
9b	Maxillule, endopod 1-segmented, with 4 or 5 setae.....	10
10a/9b	Maxillule, endopod 1-segmented, with 4 setae	11
10b	Maxillule, endopod 1-segmented, with 5 setae	12
11a/10a	Maxillule, exopod with 8 setae; coxal epipodite with 6 setae. Maxilla with strong spines	<i>Haloptilus plumosus</i>
11b	Maxillule, exopod with 11 setae; coxal epipodite with 9 setae. Maxilla without strong spines	<i>Haloptilus tenuis</i>

Available online 14 November 2022


* Corresponding author.

E-mail address: ctshih@mail.ntou.edu.tw (C.-t. Shih).



12a/10b	Antenna: endopod/exopod ratio smaller than 2. Maxillule, exopod with 11 setae. Prosome length/width ratio larger than 4	<i>Haloptilus austini</i>
12b	Antenna: endopod/exopod ratio larger than 2. Maxillule, exopod with 9 or less setae. Prosome length/width ratio 3 or less	13
13a/12b	Antenna, endopod/exopod ratio larger than 3. Maxillule exopod with 7 setae. Prosome/uosome length ratio larger than 7.5	<i>Haloptilus longicornis</i>
13b	Antenna, endopod/exopod ratio smaller than 2.5. Maxillule exopod with 9 setae. Prosome/uosome length ratio 7 or less	14
14a/13b	Maxilla with strong spines. Prosome/uosome length ratio about 6	<i>Haloptilus longicirrus</i>
14b	Maxilla without strong spines. Prosome/uosome length ratio about 7	<i>Haloptilus paralongicirrus</i>
15a/1b	Right and left legs 5, distal end of endopod not exceeding distal border of exopod segment 1	<i>Haloptilus mucronatus</i>
15b	Right and left legs 5, distal end of endopod clearly exceeding distal border of exopod segment 1	16
16a/15b	Both terminal spines of left leg 5 exopod segment 3 of equal size	<i>Haloptilus longicornis</i>
16b	Both terminal spines of left leg 5 exopod segment 3 unequal in size	17
17a/16b	Inner coxal projection present on both leg 5	18
17b	Inner coxal projection present only on left leg 5	19
18a/17a	Inner coxal projection of right leg 5 larger than that of left leg 5. Right leg 5 exopod segment 3 terminal spine shorter than exopod segment 3	<i>Haloptilus chierchiae</i>
18b	Inner coxal projection of right leg 5 smaller than that of left leg 5. Right leg 5 exopod segment 3 terminal spine longer than exopod segment 3	<i>Haloptilus paralongicirrus</i>
19a/17b	Prosome/uosome length ratio larger than 6. Caudal ramus nearly rounded	<i>Haloptilus spiniceps</i>
19b	Prosome/uosome length ratio smaller than 5. Caudal ramus squared or rectangular	20
20a/19b	Left leg 5: distal end of inner coxal projection blunt; endopod extended to distal border of exopod segment 2	<i>Haloptilus ornatus</i>
20b	Left leg 5: distal end of inner coxal projection pointed; endopod extended to midlength of exopod segment 2	21
21a/20b	Right leg 5 coxa with an inner spine; right leg 5 exopod segment 2 with an inner conical protrusion proximally. Right leg 5 terminal spine shorter than exopod segment 3	<i>Haloptilus acutifrons</i>
21b	Left leg 5 coxa without inner spine; right leg 5 exopod segment 2 without inner protrusion. Right leg 5 terminal spine equal to or longer than exopod segment 3	22
22a/21b	Antenna: endopod/exopod length ratio 4. Caudal ramus length/width ratio 1	<i>Haloptilus fertilis</i>
22b	Antenna: endopod/exopod length ratio 2. Caudal ramus length/width ratio 3	<i>Haloptilus oxycephalus</i>
22c	Antenna unknown. Caudal ramus length/width ratio 1.3	<i>Haloptilus tenuis</i>

Haloptilus acutifrons (Giesbrecht, 1892) (Size: female, 2.50–4.66 mm; male, 2.00–3.19 mm)




Sars, 1925: pl. 74. *Haloptilus acutifrons* (Giesbrecht, 1892), Female. 1/2. habitus (dorsal/lateral); 3. forehead (lateral); 4. antenna; 5. mandible; 6. maxillule; 7. maxilla; 8. maxilliped; 9. leg 1; 10. leg 3; 11. leg 5.

Bradford-Grieve, 1999: Fig. 30. Female. A/B. habitus (dorsal/lateral); C. leg 5. Male. D. habitus (dorsal); E. leg 5 (posterior, endopod incomplete).

Bradford-Grieve et al., 1999: Fig. 7.265. Female. maxillule.

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 23, 2021].

Haloptilus austini Grice, 1959 (Size: female, 3.06–3.33 mm; male, unknown)




Haloptilus austini Grice, 1959

Grice, 1959: figs. 1–18. Female. 1. habitus (dorsal); 2/3. forehead (lateral/ventral); 4/5. urosome (dorsal/lateral); 6. genital double-somite (ventral); 7. antenna; 8. mandible; 9. biting edge of mandible; 10. maxillule; 11. maxillule exopod; 12. maxilla; 13. maxilliped; 14–18. Legs 1–5

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 23, 2021].

Haloptilus chierchiai (Giesbrecht, 1889) (Size: female, 4.05–5.28 mm; male, 3.65–3.95 mm)



Sars



Sewell


Haloptilus chierchiai (Giesbrecht, 1889)

Sars, 1925: pl. 70. Female. 1/2. habitus (dorsal/lateral); 3. antennule; 4. antenna; 5/6. mandible & gnathobase; 7. maxillule; 8. maxilla; 9. maxilliped; 10. leg 1; 11. leg 3; 12. leg 5; 13. urosome (dorsal).

Sewell, 1947: Fig. 49. Female. A. mandible; C. maxillule; E. maxilla; H. leg 1. Male. B. mandible; D. maxillule; F. maxilla; G. maxilliped; I. leg 5 (posterior).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 23, 2021].

Haloptilus fertilis (Giesbrecht, 1892) (Size: female, unknown; male, 2.47–3.20 mm)




Haloptilus fertilis (Giesbrecht, 1892)

Park, 1968: pl. 11. Male. 7. habitus (dorsal); 8. urosome (dorsal); 9. left antennule; 10. antenna; 11 & 12. mandible & cutting edge; 13. maxillule; 14. maxilla; 15. maxilliped; 16–19. leg 1–4; 20. leg 5 (posterior).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 23, 2021].

Haloptilus fons Farron, 1908 (Size: female, 4.80–6.60 mm; male, unknown)




Haloptilus fons Farron, 1908


Farron, 1908: pl. 7. Female. 11. habitus (dorsal); 12. leg 5; 13. mandible cutting edge; 14. maxillule; 15. urosome (dorsal).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 14, 2021].

Haloptilus longicirrus Brodsky, 1950 (Size: female, 2.00–3.77; male, unknown)



*330-340




*341-342

Haloptilus longicirrus Brodsky, 1950


Park, 1970. Female. *330–340: 330/331. habitus (dorsal/lateral); 332/333/334. forehead (dorsal/lateral/ventral); 335/336. urosome (dorsal/lateral); 337. genital double-somite (ventral); 338. antenna; 339/340. mandible/cutting edge; *341–342: 341. maxillule; 342. leg 5 (anterior).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. Accessed May 23, 2021].


Haloptilus longicornis (Claus, 1863) (Size: female: 1.72–2.63 mm; male, 1.16–1.37 mm)



Sars



Giesbrecht



Original

Haloptilus longicornis (Claus, 1863)


Sars, 1925: pl. 82. Female. habitus (dorsal); a². Antenna; M. mandible; m. maxillule; p5. Leg 5; R. rostrum (ventral); Urs. Urosome (dorsal).

Giesbrecht, 1892: pl. 27. Male. 3. Distal portion of antennule; 29. Leg 5 (posterior).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 23, 2021].

Original. *Haloptilus longicornis* (Claus, 1863), Female, 2.06 mm, East China Sea (25°17'N, 121°35'E): a/b. habitus (dorsal/lateral); c. mandible, tip of gnathobase; d. right maxillule, posterior; e. right leg 5, posterior. Scales: a, b = 0.5 mm; c = 0.2 mm; d = 0.2 mm; e = 0.2 mm.

Haloptilus mucronatus (Claus, 1863) (Size: female, 3.00–3.60 mm; male, 2.17–2.68 mm)




Haloptilus mucronatus (Claus, 1863)

Sars, 1925: pl. 73. Female. 11. habitus (dorsal); 12. forehead (lateral); 13. mandible; 14. maxillule; 15. maxilla.
Giesbrecht, 1892: pl. 27. Male. 39. leg 5 (anterior).


Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 23, 2021].

Original. *Haloptilus mucronatus* (Claus, 1863), Female, 2.06 mm, East China Sea (25°17'N, 121°35'E): a. habitus, dorsal; b. habitus, lateral; c. mandible, tip of gnathobase.


Haloptilus ornatus (Giesbrecht, 1892) (Size: female, 3.00–5.33 mm; male, 2.75–3.05 mm.



Sars




Giesbrecht

*Haloptilus ornatus* (Giesbrecht, 1892)

Sars, 1925: pl. 73. Female. 1. habitus (dorsal); 2. forehead (Lateral); 3. mandible gnathobase; 4. maxillule; 5. maxilla.

Giesbrecht, 1892. Male. pl. 27, Fig. 38. leg 5 (posterior); pl. 42, Fig. 9. habitus (dorsal).


Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en> [Accessed May 17, 2021].

Haloptilus ornatus (Giesbrecht, 1892)

Haloptilus ornatus (Giesbrecht, 1892)

Original. Female, 3.98 mm, east of Taiwan (22°50'N, 121°55'E). a/b. habitus (dorsal/lateral); c. right mandible, gnathobase; d. right maxilla, distal end of endopod; e. leg 5.

Haloptilus oxycephalus (Giesbrecht, 1889) (Size: female, 3.00–5.20 mm; male, 2.40–2.85 mm))




Haloptilus oxycephalus (Giesbrecht, 1889)

Sars, 1925: pl. 74. Female. 12. habitus (dorsal); 13. forehead (lateral); 14. mandible; 15. maxillule; 16. maxilliped.

Park, 1988: Fig. 6. Male. a. habitus (dorsal); b/c. forehead (dorsal/ventral); d. pediger 5 & urosome (dorsal); e. left antennule; f. distal segments of left antennule; g. leg 5 (posterior).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 23, 2021].

Haloptilus paralongicirrus Park, 1970 (Size: female, 2.31–2.70 mm; male, 1.67–2.20 mm)




Haloptilus paralongicirrus Park, 1970

Park, 1970: figs. 343–355. Female. 343. habitus (dorsal); 344/345. forehead (dorsal/ventral); 346. urosome (dorsal); 347. genital double-somite (ventral); 348. antenna; 349/350. mandible & biting edge; 351. maxillule; 352. maxilla; 353. maxilliped; 354. leg 1; 355. leg 5.

Roe, 1975: Fig. 24. Male. a. posterior prosome & urosome (dorsal); b. forehead (dorsal); c. rostrum (lateral); e. antennule seg. 15–20; f. antennae; g/h. mandible/cutting edge; i. maxilla; j. maxilliped; k/l. legs 1/2; m. leg 5, endopods omitted (posterior).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 23, 2021].


Haloptilus paralongicirrus Park, 1970



Haloptilus paralongicirrus Park, 1970

Original. Female, 2.42 mm, South China Sea, 9°29'N, 114°42'E, 2438 m a/b halibut (dorsal/lateral).

Haloptilus plumosus (Claus, 1863) (Size: female, 4.15–4.20 mm; male, unknown)




Haloptilus plumosus (Claus, 1863)

Bradford-Grieve et al., 1999: Fig. 7.278. Female. A. maxillule; B. maxilla; C. maxilliped. Male. D. leg 5.

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 23, 2021].

Haloptilus spiniceps (Giesbrecht, 1892) (Size: female, 3.20–5.45 mm; male, 2.55–2.81 mm)




Haloptilus spiniceps (Giesbrecht, 1892)

Sars, 1925: pl. 73. Female. 6. habitus (dorsal); 7. forehead (lateral); 8. mandible; 9. maxillule; 10. maxilla.
 Giesbrecht, 1892: pl. 27. Male. 5. distal portion of antennule; 40. leg 5 (anterior).


Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 23, 2021].

Original. *Haloptilus spiniceps* (Giesbrecht, 1892), Female, 4.00 mm: a/b; habitus (dorsal/lateral); c. right mandible, gnathobase (posterior); d. right maxillule (anterior); e. right leg 5 (posterior). Scales: a, b = 1.00 mm; c = 0.50 mm; d = 0.50 mm; e = 0.50 mm.

Haloptilus tenuis Farron, 1908 (Size: female, 4.40–4.62 mm; male, 3.50 mm)



Sars



Bradford-Grieve

Haloptilus tenuis Farron, 1908

G.O. Sars in Résult. Camp. Scient. Prince Albert I, 69, pls.1-127 (1924). Pl.LXIX, figs.1-15. Female: 1, habitus (dorsal); 2, forehead (lateral); 3, Antennule; 4, Antenna; 5, mandible; 6, maxillule; 7, maxilla; 8, maxilliped; 9, leg 1; 10, leg 3; 11, leg 5; 12, urosome. Male: 13, forehead (lateral); 14, mandible (without palp: broken); 15, leg 5.

J.M. Bradford-Grieve, 1999. Fig.37. Female: A, habitus (dorsal); B, anterior head (lateral left side); C, maxillule; D, maxilla.

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 24, 2021].