

Volume 30  
Issue 5 Review-Special Issue (Part 2)

Article 26

## Key to the species of Bathypontiidae Temorites occurring in the China Seas

Chang-tai Shih

*National Taiwan Ocean University, ctshih@mail.ntou.edu.tw*

Qing-Chao Chen

*South China Sea Institute of Oceanology*

Yang-Chi Lan

*Fisheries Research Institute*

Shih-Hui Hsiao

*National Taipei University of Education*

Chi-Yu Weng

*Institute of Oceanography, National Taiwan University/Ocean Data Bank of the Ministry of National Science and Technology Council, Taiwan*

Follow this and additional works at: <https://jmstt.ntou.edu.tw/journal>

 Part of the Fresh Water Studies Commons, Marine Biology Commons, Ocean Engineering Commons, Oceanography Commons, and the Other Oceanography and Atmospheric Sciences and Meteorology Commons

---

### Recommended Citation

Shih, Chang-tai; Chen, Qing-Chao; Lan, Yang-Chi; Hsiao, Shih-Hui; and Weng, Chi-Yu (2022) "Key to the species of Bathypontiidae Temorites occurring in the China Seas," *Journal of Marine Science and Technology*: Vol. 30: Iss. 5, Article 26.

DOI: 10.51400/2709-6998.2617

Available at: <https://jmstt.ntou.edu.tw/journal/vol30/iss5/26>

This Review-Taxonomic Index is brought to you for free and open access by Journal of Marine Science and Technology. It has been accepted for inclusion in Journal of Marine Science and Technology by an authorized editor of Journal of Marine Science and Technology.

## REVIEW-TAXONOMIC INDEX

# Key to the Species of Bathypontiidae *Temorites* Occurring in the China Seas

Chang-tai Shih <sup>a,\*</sup>, Qing-Chao Chen <sup>b</sup>, Yang-Chi Lan <sup>c</sup>, Shih-Hui Hsiao <sup>d</sup>, Chi-Yu Weng <sup>e</sup>

<sup>a</sup> National Taiwan Ocean University, Taiwan

<sup>b</sup> South China Sea Institute of Oceanology, China

<sup>c</sup> Fisheries Research Institute, Taiwan

<sup>d</sup> National Taipei University of Education, Taiwan

<sup>e</sup> Institute of Oceanography, National Taiwan University/Ocean Data Bank of the Ministry of National Science and Technology Council, Taiwan

*Temorites brevis*(4a/11a/f)\*, *elongata*(5a/12a/f)\*, *longicornis*(12b/f)\* (female unknown), *minor*(2a/9a/f)\*, *sarsi*(6a/10a/f)\*, *similis*(3a/8b/f)\*, *spinifera*(8a/f)\*

1a	Female .....	2
1b	Male .....	7
2a/1a	Leg 5, segment 3 shorter than segment 2; segment 3 with 1 terminal spine, terminal spine more than 4 times as long as segment 3.....	<i>Temorites minor</i>
2b	Leg 5: segment 3 longer than segment 2; segment 3 with 2 terminal spines.....	3
3a/2b	Posterior corners of prosome asymmetrical, right corner slightly longer than left corner. Leg 5 segment 3: length ratio of inner/outer terminal seta nearly 10 .....	<i>Temorites similis</i>
3b	Posterior corners of prosome symmetrical. Leg 5 segment 3: length ratio of inner/outer terminal seta less than 7 .....	4
4a/3b	Total body length: 1.52–2.12 mm. Leg 5: segment 3 nearly 4 times as long as segment 2; segment 3 nearly twice as long as longer terminal spine .....	<i>Temorites brevis</i>
4b	Total body length longer than 2.5 mm. Leg 5: segment 3 less than 2 times as long as segment 2; segment 3 shorter than longer terminal spine .....	5
5a/4b	Total body length: 4.70–6.15 mm. Leg 5: longer terminal spine nearly 3 times as long as segment 3 .....	<i>Temorites elongata</i>
5b	Total length less than 3.5 mm. Leg 5: longer terminal spine about 1.5 times as long as segment 3 .....	6
6a/5b	Total body length: 2.60–3.04 mm. Leg 5: length ratio of segment 1/segment 2: 1.3; segment 3: inner terminal spine 4 times as long as outer terminal spine.....	<i>Temorites sarsi</i>
6b	Total body length: 3.30 mm. Leg 5: length ratio of segment 1/segment 2: 2.0; segment 3: inner terminal spine 8 times as long as outer terminal spine.....	<i>Temorites spinifera</i>
7a/1b	Left leg 5: Inner distal corner of basis with a plumose seta.....	8
7b	Left leg 5: Inner distal corner of basis without plumose seta.....	9
8a/7a	Left leg 5, length ratio of inner/outer terminal spines greater than 6. Right leg 5 with 2 terminal spines .....	<i>Temorites spinifera</i>
8b	Left leg 5, length ratio of inner/outer terminal spines smaller than 3. Right leg 5 with 1 terminal spine.....	<i>Temorites similis</i>
9a/7b	Right & left leg 5: terminal spine much longer than distal segment .....	<i>Temorites minor</i>
9b	Right & left leg 5: terminal spine much shorter than distal segment .....	10
10a/9b	Left leg 5, 5-segmented, inner margin of basis setose; basis of right leg 5 with an inner rounded projection on proximal half .....	<i>Temorites sarsi</i>
10b	Left leg 5, 4-segmented, inner margin of basis smooth; basis of right P5, inner margin with an inner projection, on middle section .....	11
11a/10b	Left leg 5, length ratio of left/right terminal spine: 2.5. Right leg 5: length/width ratio of segment 2 less than 2 .....	<i>Temorites brevis</i>
11b	Left leg 5, length ratio of left/right terminal spine: larger than 5. Right leg 5: length/width ratio of segment 2 larger than 12 .....	
12a/11b	Right leg 5, segment 4 straight: proximal quarter of inner margin setose .....	<i>Temorites elongata</i>
12b	Right leg 5, segment 4 curved, inner margin slightly serrated.....	<i>Temorites longicornis</i>


Available online 14 November 2022

\* Corresponding author.

E-mail address: [ctshih@mail.ntou.edu.tw](mailto:ctshih@mail.ntou.edu.tw) (C.-t. Shih).



*Temorites brevis* G.O. Sars, 1900 (Size: female, 1.52–2.12 mm; male, 1.05–1.94 mm)



Tanaka, 1965: Fig. 243. Female. a. habitus (dorsal); b. forehead (lateral); c. pediger 5 & urosome (lateral); d. mandible gnathobase; e. leg 1; f. leg 2; g. leg 3 coxa & basis; h. leg 5. Male. i. pediger 5 & urosome (lateral); j. antennule; k. leg 2; l. leg 5 (posterior).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 24, 2021].

Original. *Temorites brevis* G.O. Sars, 1900, Female, 1.78 mm, east of Taiwan (23°57'N, 122°16'E, 800 m): a/b. habitus (dorsal/lateral); c. leg 5 (posterior).

*Temorites elongata* G.O. Sars, 1905 (Size: female, 4.70–6.15 mm; male, 4.68–5.55 mm)




fig. 4





fig. 5

*Temorites elongata* G.O. Sars, 1905


Bradford-Grieve, 1999: Fig. 4. Female. A. habitus (lateral); B. antennule; C. antenna; D. mandibular blade; E. mandibular palp; F. maxillule; G. maxilla; H. maxilliped; I-M. legs 1–5. Fig. 5. Male. A. habitus (laterodorsal); B. right antennule; C. antenna; D. mandibular blade; E. mandibular palp; F. maxillule; G. maxilla; H. maxilliped; I-L. legs 1–4; M. leg 5 (posterior).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 24, 2020].

*Temorites elongata* G.O. Sars, 1905*Temorites elongata* G.O. Sars, 1905

Original. *Temorites elongata* G.O. Sars, 1905, Female, 5.64 mm, South China Sea ( $16^{\circ}40.8'N$ ,  $138^{\circ}54.0'E$  2000 m), a/b. habitus (dorsal/lateral); c. rostrum (anteroventral) d. leg 5. Male, 5.84 mm, South China Sea ( $9^{\circ}29'N$ ,  $114^{\circ}42'E$ , 2438 m), a/b. habitus (dorsal/lateral); c. leg 5 (posterior); d. tip of distal segment.

*Temorites longicornis* (Tanaka, 1965) (Size: female, Unknown; male: 5.18 mm)




*Temorites longicornis* (Tanaka, 1965)

Tanaka, 1965: Fig. 242. Male. a. habitus (dorsal); b. forehead (lateral); c. pediger 5 & urosomites 1 & 2 (lateral); d. rostrum (dorsal); e. right antennule; f. mandible gnathobase; g-i. legs 1–3; j. leg 5 (anterior).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 25, 2021].

*Temorites minor* (Wolfenden, 1906) (Size: female, 2.50–2.80 mm; male, 2.30–2.51 mm)




*Temorites minor* (Wolfenden, 1906)

Deevey, 1973: Fig. 3. Female. a/b. habitus (dorsal/lateral); e. leg 5. Male. d. leg 5 (anterior); h/i. habitus (dorsal/lateral).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 25, 2020].

*Temorites sarsi* (Grice & Hülsemann, 1967) (Size: female, 2.60–3.04 mm; male, 2.74–3.04 mm)




*Temorites sarsi* (Grice & Hülsemann, 1967)

Deevey, 1967: Fig. 8. Female. a/b. habitus (lateral/dorsal); c. antenna; d. maxillule; e. mandible; f. maxiliped; g. syncoxa of maxilla; h. leg 5; k. mandible gnathobase. Male. a. habitus (lateral); i. leg 5 (anterior).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 25, 2021].

*Temorites similis* (Tanaka, 1965) (Size: female, 2.60–3.12 mm; male, unknown)




Tanaka, 1965: Fig. 241. Female. a. habitus (dorsal); b. forehead (lateral); c. rostrum (frontal); d. pediger 5 & genital double-somite (lateral); e. posterior corner of prosome (left lateral); f. mandible; g. maxilla; h. leg 1; i. leg 5. Male. j. pediger 5 & urosome (dorsal); k. leg 5 (posterior).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 25, 2021].

Original. *Temorites similis* (Tanaka, 1965), Male, 2.80 mm., South China Sea, 6°46'N, 112°43'E, 1200 m. a/b. habitus (dorsal/lateral); c. rostrum (anteroventral); leg 5 (anterior).

*Temorites spinifera* (A. Scott, 1909) (Size: female, 3.30 mm; male, 2.60–3.00 mm)



*Temorites spinifera* (A. Scott, 1909)

Scott, 1909: pl. 3. Female. 3. urosome (dorsal); 4. antennule; 6. antenna; 7. mandible; 8. maxillule; 9. maxilla; 10. maxilliped; 11. leg 1; 12. leg 2; 14. leg 4; 16. leg 5. Male. 1. habitus (dorsal); 2. pedigers 4/5 & urosomite 1 (lateral); 5. right antennule; 13. leg 2; 15. leg 5 (posterior).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 25, 2021].