



Key to the species of Calanidae Neocalanus occurring in the China Seas

Chang-tai Shih

National Taiwan Ocean University, ctshih@mail.ntou.edu.tw

Qing-Chao Chen

South China Sea Institute of Oceanology

Yang-Chi Lan

Fisheries Research Institute

Shih-Hui Hsiao

National Taipei University of Education

Chi-Yu Weng

Institute of Oceanography, National Taiwan University/Ocean Data Bank of the Ministry of National Science and Technology Council, Taiwan

Follow this and additional works at: <https://jmstt.ntou.edu.tw/journal>



Part of the [Fresh Water Studies Commons](#), [Marine Biology Commons](#), [Ocean Engineering Commons](#), [Oceanography Commons](#), and the [Other Oceanography and Atmospheric Sciences and Meteorology Commons](#)

Recommended Citation

Shih, Chang-tai; Chen, Qing-Chao; Lan, Yang-Chi; Hsiao, Shih-Hui; and Weng, Chi-Yu (2022) "Key to the species of Calanidae Neocalanus occurring in the China Seas," *Journal of Marine Science and Technology*: Vol. 30: Iss. 5, Article 31.

DOI: 10.51400/2709-6998.2622

Available at: <https://jmstt.ntou.edu.tw/journal/vol30/iss5/31>

This Review-Taxonomic Index is brought to you for free and open access by Journal of Marine Science and Technology. It has been accepted for inclusion in Journal of Marine Science and Technology by an authorized editor of Journal of Marine Science and Technology.

REVIEW-TAXONOMIC INDEX

Key to the Species of Calanidae *Neocalanus* Occurring in the China Seas

Chang-tai Shih ^{a,*}, Qing-Chao Chen ^b, Yang-Chi Lan ^c, Shih-Hui Hsiao ^d, Chi-Yu Weng ^e

^a National Taiwan Ocean University, Taiwan

^b South China Sea Institute of Oceanology, China

^c Fisheries Research Institute, Taiwan

^d National Taipei University of Education, Taiwan

^e Institute of Oceanography, National Taiwan University/Ocean Data Bank of the Ministry of National Science and Technology Council, Taiwan

Neocalanus cristatus (2a/4a/f), *gracilis* (3a/5a/f), *robustior* (3b/5b/f)

1a	Female	2
1b	Male	4
2a/1a	Anterior head with a shallow crest	<i>Neocalanus cristatus</i>
2b	Anterior head without crest	3
3a/2b	Antennule, when bent posteriorly, distal 6 segments exceeding distal margin of caudal rami. Leg 1: process at distal outer corner of endopod segment 1 insignificant; endopod segment 2 distal outer margin with an upward hooked process	<i>Neocalanus gracilis</i>
3b	Antennule, when bent posteriorly, distal 8 segments exceeding distal margin of caudal rami. Leg 1: process at distal outer corner of endopod segment 1 distinct; endopod segment 2 distal outer margin without hooked process	<i>Neocalanus robustior</i>
4a/1b	Anterior head with crest. Left leg 5: exopod 2-segmented. Total length longer than 5 mm	<i>Neocalanus cristatus</i>
4b	Anterior head without crest. Left leg 5: exopod 3-segmented. Total length shorter than 4 mm	5
5a/4b	Left Leg 5, endopod reaching or slightly exceeding distal border of exopod segment 1. Right leg 5, exopod reaching distal border	<i>Neocalanus gracilis</i>
5b	Left leg 5, endopod reaching midlength of exopod segment 1. Right exopod, nearly reaching midlength of left exopod segment 1	<i>Neocalanus robustior</i>

Available online 14 November 2022

* Corresponding author.

E-mail address: ctshih@mail.ntou.edu.tw (C.-t. Shih).



Neocalanus cristatus (Kröyer, 1848) (Size: female, 7.60–10.40 mm; male, 6.76–9.60 mm)




Fig. 77





Fig.78

Neocalanus cristatus (Kröyer, 1848)

Brodsky et al., 1983: Fig. 77. Female. habitus (dorsal); abd. urosome (lateral/ventral); abd 1. genital double-somite; ce. forehead (lateral); Gn. genital field; mx2. maxilla; p1/4. Leg 1/4; p5, leg 5. Fig. 78. Male. habitus (dorsal); Ce. forehead (lateral); p4. leg 4; p5. leg 5 (posterior); P5 ex 2 + 3. Right leg 5, exopod segment 2&3 (anterior).


Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 27, 2021].

Neocalanus gracilis (Dana, 1852) (size: female, 1.80–4.44 mm; male, 1.60–3.40 mm)



Chen & Zhang, 1965


Neocalanus gracilis (Dana, 1852)



Bradford-Grieve et al., 1999

Chen & Zhang, 1965: pl. 1. *Neocalanus gracilis* (Dana, 1852), Female. 8/9. habitus (dorsal/lateral), 10. leg 1. Bradford-Grieve et al., 1999: Fig. 720. Male. habitus (dorsal); leg 5 (anterior).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 27, 2021].

Neocalanus gracilis (Dana, 1852)


Female

Neocalanus gracilis (Dana, 1852)


Male

Original. *Neocalanus gracilis* (Dana, 1852), **Female**, 3.16 mm, East China Sea (25°25'N 122°15'E): a/b. habitus (dorsal/lateral). **Male**, 2.48 mm, East China Sea (25°25'N 122°15'W): a/b. habitus (dorsal/lateral); c. endopod of right leg 1, posterior; d. exopod segment 3 of right leg 1, posterior; e. leg 5, posterior.

Neocalanus robustior (Giesbrecht, 1888) (Size: female, 3.00–4.65 mm; male, 2.52–3.60 mm)




Chen & Zhang, 1965



Chen et al., 1974

Neocalanus robustior (Giesbrecht, 1888)



Original

Chen & Zhang, 1965: pl. 2. *Neocalanus robustior* (Giesbrecht, 1888), Female. 1/2. habitus (dorsal/lateral); 3. leg 1 (anterior).

Chen et al., 1974: pl. 1. Male. 4. habitus (dorsal); 5. left leg 5 (posterior).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 27, 2021].

Original. *Neocalanus robustior* (Giesbrecht, 1888), Female, 3.44 mm, East of Taiwan (23°55.0'N 122°25.1'E): a/ b. habitus (dorsal/lateral); c. part of leg 1.