



Key to the species of Clausocalanidae Clausocalanus occurring in the China Seas

Chang-tai Shih
National Taiwan Ocean University, ctshih@mail.ntou.edu.tw

Qing-Chao Chen
South China Sea Institute of Oceanology

Yang-Chi Lan
Fisheries Research Institute

Shih-Hui Hsiao
National Taipei University of Education

Chi-Yu Weng
Institute of Oceanography, National Taiwan University/Ocean Data Bank of the Ministry of National Science and Technology Council, Taiwan

Follow this and additional works at: <https://jmstt.ntou.edu.tw/journal>



Part of the [Fresh Water Studies Commons](#), [Marine Biology Commons](#), [Ocean Engineering Commons](#), [Oceanography Commons](#), and the [Other Oceanography and Atmospheric Sciences and Meteorology Commons](#)

Recommended Citation

Shih, Chang-tai; Chen, Qing-Chao; Lan, Yang-Chi; Hsiao, Shih-Hui; and Weng, Chi-Yu (2022) "Key to the species of Clausocalanidae Clausocalanus occurring in the China Seas," *Journal of Marine Science and Technology*. Vol. 30: Iss. 5, Article 36.

DOI: 10.51400/2709-6998.2627

Available at: <https://jmstt.ntou.edu.tw/journal/vol30/iss5/36>

This Review-Taxonomic Index is brought to you for free and open access by Journal of Marine Science and Technology. It has been accepted for inclusion in Journal of Marine Science and Technology by an authorized editor of Journal of Marine Science and Technology.

REVIEW-TAXONOMIC INDEX

Key to the Species of Clausocalanidae *Clausocalanus* Occurring in the China Seas (*Modified and Expanded from Frost & Fleminger, 1968)

Chang-tai Shih ^{a,*}, Qing-Chao Chen ^b, Yang-Chi Lan ^c, Shih-Hui Hsiao ^d, Chi-Yu Weng ^e

^a National Taiwan Ocean University, Taiwan

^b South China Sea Institute of Oceanology, China

^c Fisheries Research Institute, Taiwan

^d National Taipei University of Education, Taiwan

^e Institute of Oceanography, National Taiwan University/Ocean Data Bank of the Ministry of National Science and Technology Council, Taiwan

Clausocalanus arcuicornis, *dubius*(13a/f) (male only), *farrani*(4a/18b/f), *furcatus*(2a/19a/f), *ingens*(7a/22a/f), *jobei*(4b/17a/f), *laticeps*(6a/21a/f), *lividus*(8a/23a/f), *mastigophorus*(8b/24a/f), *minor*(10a/18a/f), *parapergens*(12b/24b/f), *paululus*(9a/15a/f), *pergens*(12a/20a/f)

1a	Female	2
1b	Male	13
2a/1a	Genital double-somite and urosomite 3 subequal in length.....	<i>Clausocalanus furcatus</i>
2b	Genital double-somite more than 1.5 times as long as urosomite 3	3
3a/2b	In lateral view, ventral profile of genital double-somite conspicuously protuberant anterior to genital pores	4
3b	In lateral view, ventral profile of genital double-somite not conspicuously protuberant anterior to genital pores.....	5
4a/3a	In lateral view, rostrum short, thick, and straight.....	<i>Clausocalanus farrani</i>
4b	In lateral view, rostrum long, slender, and curved	<i>Clausocalanus jobei</i>
5a/3b	Leg 5, segment 3 less than 2 times as long as segment 1.....	6
5b	Leg 5, segment 3 more than 2 times as long as segment 1.....	9
6a/5a	Front of head usually strongly vaulted. In leg 3, distance between basipodal spiniform processes 2 and 3 equal to or more than 2 times the proximal width of process 3, processes 2 and 3 uniformly tapered from base. Caudal ramus more than 1.65 times as long as wide.....	<i>Clausocalanus laticeps</i>
6b	Front of head not vaulted. In leg 3, distance between basipodal spiniform processes 2 and 3 less than 1.5 times the proximal width of process 3, processes 2 and 3 not uniformly tapered from base. Caudal ramus less than 1.65 times as long as wide	7
7a/6b	In lateral view, rostrum slender and curved, front of head broadly rounded, protuberant above rostrum. Greatest anterior extension of forehead well dorsal to frontal organ	<i>Clausocalanus ingens</i>
7b	In lateral view, rostrum thick, straight or curved, front of head broadly rounded but not markedly protuberant above rostrum. Greatest anterior extension of forehead at the level of frontal organ, or, rarely, slightly dorsal to the frontal organ	8
8a/7b	Rostrum usually straight in lateral view, tapered uniformly in either lateral or anteroventral views	<i>Clausocalanus lividus</i>
8b	Rostrum usually curved ventroposteriorly in lateral view, not tapered uniformly in either lateral or anteroventral views	<i>Clausocalanus mastigophorus</i>
9a/5b	In leg 3, distance between basipodal spiniform processes 2 and 3 more than 2 times proximal width of process 3	<i>Clausocalanus paululus</i>
9b	In leg 3, distance between basipodal spiniform processes 2 and 3 less than 1.5 times proximal width of process 3.....	10
10a/9b	In lateral view, ventral profile of genital double-somite with a prominent step posterior to genital pore	<i>Clausocalanus minor</i>

Available online 14 November 2022


* Corresponding author.

E-mail address: ctshih@mail.ntou.edu.tw (C.-t. Shih).



10b	In lateral view, ventral profile of genital double-somite without prominent step posterior to genital pore	11
11a/10b	In lateral view, dorsal lobe of seminal receptacle, not bulb-shaped, arising from anterior edge of ventral lobe	<i>Clausocalanus arcuicornis</i>
11b	In lateral view, dorsal lobe of seminal receptacle, usually bulb-shaped, portion of ventral lobe visible anterior to dorsal lobe	12
12a/11b	In lateral view, dorsal lobe of seminal receptacle digital form or slightly constricted in region of attachment to ventral lobe of seminal receptacle. In lateral view, posterolateral corner of prosome somewhat rounded. Antennule, segment 2 less than 0.95 times as long as segment 24. In leg 3, exopod segment 3 less than 1.20 times as long as terminal spine. In leg 4, exopod segment 3 less than 1.30 times as long as terminal spine	<i>Clausocalanus pergens</i>
12b	In lateral view, dorsal lobe of seminal receptacle bulb-shaped constricted in region of attachment to ventral lobe of seminal receptacle. In lateral view, posterolateral corner of prosome somewhat angular. Antennule, segment 2 about 1.07–1.26 times as long as segment 24. In leg 3, exopod segment 3: 1.40–1.70 times as long as terminal spine. In leg 4, exopod segment 3: 1.40–1.60 times as long as terminal spine	<i>Clausocalanus parapergens</i>
13a/1b	Right leg 5 extended passing distal margin of left leg 5 segment 1. Urosomites 2, 3, and 4 subequal in length	<i>Clausocalanus dubius</i>
13b	Right leg 5 extended not passing mid length of left leg 5 segment 1. Urosomites 2 at least 1.5 times as long as urosomite 3	14
14a/13b	In lateral view, seminal vesicle extending within prosome anterior to level of leg 1 articulation with body. Spermatophore extending by at least one-quarter of its length anterior to articulation of pedigers 1 and 2	15
14b	In lateral view, seminal vesicle not extending within prosome anterior to level of leg 1 articulation with body. Spermatophore not extending or extending by less than one-eighth of its length anterior to articulation of pedigers 1 and 2	19
15a/14a	Total length less than 0.70 mm	<i>Clausocalanus paululus</i>
15b	Total length greater than 0.70 mm	16
16a/15b	Prosome less than 5.7 times as long as urosomite 2	<i>Clausocalanus arcuicornis</i>
16b	Prosome more than 5.7 times as long as urosomite 2	17
17a/16b	Left leg 5, armature of segment 5 including 2 thick, curved, spiniform setae; segment 2 length more than 1.45 times as great as width of urosomite 2	<i>Clausocalanus jobei</i>
17b	Left leg 5, armature of segment 5 including slender, usually straight, setae; segment 2 length less than 1.45 times as great as width of urosomite 2	18
18a/17b	Right leg 5, 2- or 3-segmented, distal segments large. Antennule, fused segments 20/21 more than 1.35 times as long as segment 22	<i>Clausocalanus minor</i>
18b	Right leg 5, always 2-segmented, distal segments reduced. Antennule, fused segments 20/21 less than 1.35 times as long as segment 22	<i>Clausocalanus farrani</i>
19a/14b	In lateral view, rostrum not knoblike and not protruding ventrally. Longer ramus of leg 5 usually on right side	<i>Clausocalanus furcatus</i>
19b	In lateral view, rostrum knoblike and protruding ventrally. Longer ramus of leg 5 usually on left side	20
20a/19b	Total length less than 0.80 mm	<i>Clausocalanus pergens</i>
20b	Total length greater than 0.80 mm	21
21a/20b	In leg 3, spiniform processes 1–3 of basis slender and tapered uniformly from base. Caudal ramus more than 1.6 times as long as wide	<i>Clausocalanus laticeps</i>
21b	In leg 3, spiniform processes 2 and 3 of basis usually stout and not tapered uniformly from base. Caudal ramus less than 1.6 times as long as wide	22
22a/21b	In left leg 5, segment 2 less than 4.4 times as long as wide	<i>Clausocalanus ingens</i>
22b	In left leg 5, segment 2 more than 4.4 times as long as wide	23
23a/22b	Urosomite 2 more than 1.35 times as long as leg 5 segment 2	<i>Clausocalanus lividus</i>
23b	Urosomite 2 less than 1.35 times as long as leg 5 segment 2	24
24a/23b	Prosome more than 6.45 times as long as urosomite 2. Right leg 5, two distal segments usually well developed	<i>Clausocalanus mastigophorus</i>
24b	Prosome less than 6.45 times as long as urosomite 2. Right leg 5, two distal segments usually reduced	<i>Clausocalanus parapergens</i>

Clausocalanus arcuicornis (Dana, 1849) (Size: Female, 1.08–1.62 mm; male, 0.90–1.25 mm)




Female

Male

Clausocalanus arcuicornis (Dana, 1849)


Frost & Fleminger, 1968: pls. 29–31. *Clausocalanus arcuicornis* (Dana, 1849), Female. Pl. 29. 1a/b. habitus (lateral/dorsal). Pl. 30. 2a-d. rostrum (lateral); 2e-f. rostrum (anteroventral); 2g-i. posterior prosome & urosome (lateral); 2j. genital double-somite (ventral); 2k. leg 2 basis; 2l. leg 3 basis. Pl. 31. 3a. pediger 5 & urosome (lateral); 3b. leg 3 exopod segment 3; 3c/d. leg 5 (2 different specimens). Male. Pl. 32. 1a/b. habitus (lateral/dorsal); 1c. posterior prosome & urosome (lateral). Pl. 33. 2a. forehead (lateral); 2b. urosome (dorsal); 2c/d. leg 2 basis; 2e/f. leg 3 basis; 2g/h. leg 5 (posterior).

Razouls C. Desreumaux N., Kouwenberg J. and de Bovée F., 2005–2020. – Biodiversity of Marine Planktonic Copepods (morphology, geographical distribution and biological data). Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 03, 2021].

Clausocalanus arcuicornis (Dana, 1849)*Clausocalanus arcuicornis* (Dana, 1849)

Original. *Clausocalanus arcuicornis* (Dana, 1849), Male, 1.08 mm, Taiwan Strait (22°49'N, 120°12'E): a/b. habitus (dorsal/lateral); c. basis of right leg 2 (posterior); d. basis of left leg 3 (posterior); e. leg 5 (posterior). Scales: a, b = 0.5 mm, c, d = 0.25 mm; e = 0.25 mm.


Clausocalanus dubius Brodsky, 1950 (Size: Female, 1.80 mm; male, 0.97–1.80 mm)



Clausocalanus dubius Brodsky, 1950


Brodsky, 1950: Fig. 41. *lausocalanus dubius* Brodsky, 1950, Male habitus (dorsal); RLa. rostrum (lateral); p. rostrum (frontal); S1. Leg 1; S3B2. Leg 3, coxa & basis; S5. Leg 5.

Razouls C. Desreumaux N., Kouwenberg J. and de Bovée F., 2005–2020. – Biodiversity of Marine Planktonic Copepods (morphology, geographical distribution and biological data). Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 03, 2021].


Clausocalanus farrani Sewell, 1929 (Size: Female, 1.04–1.26 mm; male, 0.87–0.99 mm)*Clausocalanus farrani* Sewell, 1929

Frost & Fleminger, 1968: pls. 34–38. *Clausocalanus farrani* Sewell, 1929, **Female**. Pl. 34, 1a/b habitus (lateral/dorsal). 1c/d. rostrum (lateral); 1e. rostrum (anteroventral); 1f/g. forehead (lateral). Pl. 35, 2a-c. forehead (lateral); 2d. posterior prosome & genital double-somite (lateral); 2e. genital double-somite (lateral/ventral). 2f. genital single-somite (ventral). Pl. 36, 3a. posterior prosome & urosome (lateral); 3b. leg 2 basis; 3c/d. leg 3 basis; 3e. leg 3 exopod segment 3; 3f/g leg 5 (from 2 specimens). **Male**. Pl. 37, 1a/b. habitus (lateral/dorsal); 1c. forehead (lateral); 1d. posterior prosome & urosome (lateral); 1e. urosome (dorsal). Pl. 38, 2a. leg 2 basis; 2b. leg 3 basis; 2c/d. leg 5 (posterior/lateral); 2e/f. right leg 5 (anterior/lateral); 2g. left leg 5 segments 4 & 5 (posterior).

Razouls C. Desreumaux N., Kouwenberg J. and de Bovée F., 2005–2020. – Biodiversity of Marine Planktonic Copepods (morphology, geographical distribution and biological data). Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 03, 2021].

Clausocalanus farrani Sewell, 1929*Clausocalanus farrani* Sewell, 1929

Original. *Clausocalanus farrani* Sewell, 1929, Female, 1.17 mm, East China Sea (24°18'N, 121°46'E, surface): a/b. habitus (dorsal/lateral); c. basis of leg 2 (posterior); d. basis of leg 3 (posterior); e. leg 5 (posterior). Scales: a, b = 0.5 mm; c-e = 0.2 mm.

Clausocalanus furcatus (Brady, 1883) (Size: Female, 0.80–1.75 mm; male, 0.70–1.14 mm)


Female

Male

Clausocalanus furcatus (Brady, 1883)


Frost & Fleminger, 1968: pls 64–67. *Clausocalanus furcatus* (Brady, 1883), Female. Pl. 64. 1a/b. habitus (lateral/dorsal); 1c/d. rostrum (lateral/anteroventral); 1e. urosome (dorsal); 1f. right caudal ramus (dorsal); 1g, leg 2 basis; 1h. leg 3 basis. Pl. 65. 2a. posterior prosome & urosome with spermatophore attached (lateral); 2b/c. genital double-somite (lateral/ventral); 2d/e. leg 5. Male. Pl. 66. 1a/b. habitus (lateral/dorsal); 1c-e, forehead (lateral); 1f. urosome (dorsal); 1g. leg 2 basis; 1h. leg 3 basis. Pl. 67. 2 a/b, posterior prosome to urosomite 3 (dorsal); 2c. posterior prosome & urosome (lateral); 2d. leg 5 (left posterolateral); 2e/f, left/right leg 5 (anterior); 2g. right leg 5.

Razouls C. Desreumaux N., Kouwenberg J. and de Bovée F., 2005–2020. – Biodiversity of Marine Planktonic Copepods (morphology, geographical distribution and biological data). Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 03, 2020].

Clausocalanus furcatus (Brady, 1883)*Clausocalanus furcatus* (Brady, 1883)

Original. *Clausocalanus furcatus* (Brady, 1883), Female, 1.32 mm, East China Sea (25°24'N, 122°19'E, 200 m): a/b. habitus, (dorsal/lateral); c. rostrum (ventral); d. leg 2 basis (posterior); e. leg 3 basis (posterior); f. leg 5 (posterior). Scales: a, b = 0.5 mm; c-f = 0.1 mm.

Clausocalanus ingens Frost & Fleminger, 1968 (Size: Female, 1.44–1.90 mm; male, 0.99–1.08 mm)




Clausocalanus ingens Frost & Fleminger, 1968

Frost & Fleminger, 1968: pls. 19–22. *Clausocalanus ingens* Frost & Fleminger, 1968, Female. pl. 19. 1a/b. habitus (lateral/dorsal). Pl. 20. 2a. forehead (lateral); 2b/c. rostrum (lateral/anteroventral); 2d. pedigers 4–5 & genital double-somite (lateral); 2e. genital double-somite (ventral). Pl. 21. 3a. pedigers 4/5 & urosome with spermatophore attached (lateral); 3b. leg 2 basis; 3c. leg 3 basis; 3d/e. leg 5. Male. Pl. 22. 1a/b. habitus (lateral/dorsal); 1c. forehead (lateral); 1d, urosome (dorsal); 1e, urosomite 4 to caudal rami (dorsal). Pl. 23. 2a. posterior prosome & urosome (lateral); 2b. leg 2 basis; 2c. leg 3 basis; 2d/e. leg 5 (posterior/anterolateral).

Razouls C. Desreumaus N., Kouwenberg J. and de Bovée F., 2005–2020. – Biodiversity of Marine Planktonic Copepods (morphology, geographical distribution and biological data). Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 03, 2021].

Clausocalanus jobei Frost & Fleminger, 1968 (Size: Female, 1.01–1.56 mm; male, 0.87–1.07 mm)




Female

Male

Clausocalanus jobei Frost & Fleminger, 1968

Frost & Fleminger, 1968: pls.39-42. *Clausocalanus jobei* Frost & Fleminger, 1968, Female. Pl. 39. 1a/b. habitus (lateral/dorsal); 1c/d. rostrum (lateral/anteroventral); 1e. forehead (lateral); 1f. leg 2 basis; 1g. leg 3 basis. Pl. 40. 2a. posterior prosome & genital double-somite (lateral); 2b. posterior prosome & urosome with spermatophore attached (lateral); 2c-f. leg 5. Male. Pl.41. 1a/b. habitus (lateral/dorsal); 1c. forehead (lateral); 1d. posterior prosome & urosome (lateral); 1e. urosome (dorsal). Pl. 42. 2a. leg 2 basis; 2b. leg 3 basis; 2c/d. leg 5 (posterior/lateral); 2e, right leg 5 (posterior); 2f. left leg 5 segments 4/5 (posterior).

Razouls C. Desreumaus N., Kouwenberg J. and de Bovée F., 2005–2020. – Biodiversity of Marine Planktonic Copepods (morphology, geographical distribution and biological data). Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 03, 2021].

Clausocalanus laticeps Farran, 1929 (Size: Female, 1.25–1.67 mm; male, 1.01–1.10 mm)


Female

Male

Clausocalanus laticeps Farran, 1929

Frost & Fleminger, 1968: pls. 24–28. *Clausocalanus laticeps* Farran, 1929, Female. Pl. 24. 1a/b. habitus (lateral/dorsal). Pl. 25. 2a. posterior prosome & urosome with spermatophore attached (lateral); 2b. leg 2 basis; 2c–e, leg 3 basis; 2 f. leg 5. Pl. 26. 3a/b, forehead (lateral); 3c–f. rostrum (lateral); 3g. posterior prosome & genital double-somite (lateral); 3h. genital double-somite (ventral). Male. Pl. 27. 1a/b. habitus (lateral/dorsal); 1c. urosome (dorsal). Pl. 28. 2a. forehead (lateral); 2b. urosomites 4/5 & caudal rami (dorsal); 2 c/d. leg 2 basis (right and left legs of same specimen); 2e/f. leg 3 basis (right and left legs of same specimen); 2g. leg 5 (lateral).

Razouls C. Desreumaux N., Kouwenberg J. and de Bovée F., 2005–2020. – Biodiversity of Marine Planktonic Copepods (morphology, geographical distribution and biological data). Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 03, 2021].

Clausocalanus lividus Frost & Fleminger, 1968 (Size: Female, 1.22–1.98 mm; male, 1.09–1.45 mm)


Female

Male

Clausocalanus lividus Frost & Fleminger, 1968

Frost & Fleminger, 1968: pls. 14–18. *Clausocalanus lividus* Frost & Fleminger, 1968, Female. Pl. 14. 1a/b habitus (lateral/dorsal). Pl. 15. 2a/b. forehead (lateral); 2c-f. rostrum (lateral); 2g. rostrum (anterior); 2h-j. posterior prosome & genital double-somite (lateral). Pl. 16. 3a. posterior prosome & genital double-somite (lateral); 3b. genital double-somite (ventral); 3c, posterior prosome & urosome with spermatophore attached (lateral); 3d. leg 2 basis; 3e. leg 3 basis; 3f, leg 5. Male. Pl. 17. 1a/b. habitus (lateral/dorsal); 1c. forehead (lateral); 1d. urosomite 4 to caudal rami (armature incomplete) (dorsal). Pl. 18. 2a. posterior prosome & urosome (lateral); 2b. urosome (armature of caudal rami incomplete) (dorsal); 2c. leg 2 basis; 2d. leg 3 basis; 2e. leg 5 (lateral); 2f/g. right leg 5 (posterior/lateral).

Razouls C. Desreumaux N., Kouwenberg J. and de Bovée F., 2005–2020. – Biodiversity of Marine Planktonic Copepods (morphology, geographical distribution and biological data). Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 03, 2021].


Clausocalanus mastigophorus (Claus, 1863) (Size: Female, 1.17–1.94 mm; male, 1.05–1.50 mm)

Female
Clausocalanus mastigophorus (Claus, 1863)

Male

Frost & Fleminger, 1968: pls. 9–13. *Clausocalanus mastigophorus* (Claus, 1863), Female. Pl. 9. 1a/b. habitus (lateral/dorsal). Pl. 10. 2 a/b. forehead (lateral); 2c-e. rostrum (lateral); 2f. rostrum (anteroventral); 2g. leg 2 basis; 2h. leg 3 basis; 2i/j. leg 5. Pl. 11. 3a. urosome with spermatophore attached (lateral); 3b-d. posterior prosome & genital double-somite (lateral); 3e. genital-double somite (ventral). Male. Pl.12. 1a/b. habitus (lateral/dorsal); 1c. frontal region (lateral); 1d. urosome (dorsal); 1e. leg 2 basis; 1f. leg 3 basis. Pl. 13. 2a. posterior prosome & urosome (lateral); 2b. urosomite 4 to caudal rami (dorsal view); 2c. leg 5; 2d. right leg 5.

Razouls C. Desreumaux N., Kouwenberg J. and de Bovée F., 2005–2020. – Biodiversity of Marine Planktonic Copepods (morphology, geographical distribution and biological data). Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 03, 2021].


Clausocalanus mastigophorus (Claus, 1863)

Female

Clausocalanus mastigophorus (Claus, 1863)


Male

Original. *Clausocalanus mastigophorus* (Claus, 1863), female, 1.62mm, Taiwan Strait (22°49'N, 120°12'E, surface), habitus (dorsal/lateral); rostral filament; leg 2/3, basis; leg 5. Male, 1.38 mm, same locality; habitus (dorsal/lateral); leg 2/3, basis; leg 5.


Clausocalanus minor Sewell, 1929 (Size: Female, 0.94–1.32 mm; male, 0.79–1.06 mm)*Clausocalanus minor* Sewell, 1929

Frost & Fleminger, 1968: pls. 43–46. *Clausocalanus minor* Sewell, 1929, Female. Pl. 43). 1a/b. habitus (lateral/dorsal); 1c/d. rostrum (lateral/anteroventral); 1e. forehead (lateral); 1f. posterior prosome & genital double-somite (lateral); 1g. ventral half of genital double-somite (lateral). Pl. 44. 2a. genital double-somite (ventral); 2b. posterior prosome & urosome with spermatophore attached (lateral); 2c. leg 2 basis; 2d. leg 3 basis; 2e. leg 3 exopod segment 3; 2f/g. leg 5. Male. Pl. 45. 1a/b. habitus (lateral/dorsal); 1c. forehead (lateral); 1d. posterior prosome & urosome (lateral); 1e. urosome (dorsal); 1f. leg 2 basis; 1g. leg 3 basis. Pl. 46. 2a/b. leg 5 (lateral/posterior); 2c-f. right leg 5; 2g. left leg segments 4–5.

Razouls C. Desreumaux N., Kouwenberg J. and de Bovée F., 2005–2020. – Biodiversity of Marine Planktonic Copepods (morphology, geographical distribution and biological data). Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 03, 2021].

Clausocalanus minor Sewell, 1929

Female




Male

Clausocalanus minor Sewell, 1929

Original. *Clausocalanus minor* Sewell, 1929. Female, 1.15 mm, East of Taiwan (off Zhibeng, 24°18'N, 121°46'E), habitus (dorsal/lateral); rostrum; basis of legs 2 & 3; LEG 5. Male, 1.08 mm, same locality, habitus (dorsal/lateral); basis of legs 2 and 3; leg 5.

Clausocalanus parapergens Frost & Fleminger, 1968 (Size: Female, 0.97–1.38 mm; male, 0.97–1.15 mm)




Female

Male

Clausocalanus parapergens Frost & Fleminger, 1968


Frost & Fleminger, 1968: pls. 60–63. Female, Pl. 60. 1a/b, habitus (dorsal/lateral); 1c/d, rostrum (lateral/anteroventral); 1e, forehead (lateral); 1f, posterior prosome & genital double-somite (lateral); 1g, genital double-somite (ventral). Pl. 61. 2a, posterior prosome & urosome (lateral); 2b, posterior prosome & urosome with spermatophore attached (lateral); 2c, leg 2, basis; 2d, leg 3 basis; 2e, leg 3 exopod segment 3; 2f, leg 4 exopod segment 3; 2g/h, leg 5. Male. Pl. 62. 1a/b, habitus (lateral/dorsal); 1c, forehead (lateral); 1d, urosome (dorsal); 1e, leg 2 basis; 1f, leg 3 basis. Pl. 63. 2a, posterior prosome & urosome (lateral); 2b, urosomite 4/5 & caudal rami (armature incomplete) (dorsal); 2c/d, leg 5 (lateral); 2e, leg 5 (posterior); 2f, right leg 5.

Razouls C. Desreumaux N., Kouwenberg J. and de Bovée F., 2005–2020. – Biodiversity of Marine Planktonic Copepods (morphology, geographical distribution and biological data). Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 03, 2021].

Clausocalanus paululus Farran, 1926 (Size: Female, 0.65–0.86 mm; male, 0.47–0.60 mm)*Clausocalanus paululus* Farran, 1926

Frost & Fleminger, 1968: pls. 47–50. **Female.** Pl. 47. 1a/b. habitus (lateral/dorsal); 1c. rostrum (lateral). Pl. 48. 2a. posterior prosome & genital double-somite (lateral); 2b/c. genital double-somite; 2d. ventral half of genital double-somite (lateral); 2 e/f. leg 2 basis; 2g-i, leg 5 basis. Pl. 48. 3 a/b, posterior prosome & urosome with spermatophore attached (right lateral; 3c/d. leg 5 (from different specimens). **Male.** Pl. 49. 1e/f, habitus (lateral/dorsal). Pl. 50. 2a, forehead (lateral); 2b, posterior prosome & urosome (lateral); 2c, urosome (dorsal); 2d, leg 2, basis; 2e, leg 4 basis; 2f, leg 5 (anterior).

Razouls C. Desreumaus N., Kouwenberg J. and de Bovée F., 2005–2020. – Biodiversity of Marine Planktonic Copepods (morphology, geographical distribution and biological data). Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 03, 2021].

Clausocalanus pergens Farran, 1926 (Size: Female, 0.70–1.10 mm; male, 0.52–0.67 mm)*Clausocalanus pergens* Farran, 1926

Frost & Fleminger, 1968: pls. 51–55. *Clausocalanus pergens* Farran, 1926, **Female**, Pl. 51. 1a/b. habitus (lateral/dorsal); 1c. rostrum (lateral). Pl. 52. 2a. posterior prosome & genital double-somite (lateral); 2b. genital double-somite (ventral); 2c/d. leg 2 basis; 2e/f, leg 3 basis; 2g, leg 2 terminal spine; 2h, leg 3 terminal spine; 2i, leg 3 exopod segment 3; 2j, leg 4 exopod segment 3. Pl. 53. 3a, posterior prosome & urosome with spermatophore attached (lateral view); 3b, pediger 2 to urosome (lateral); 3c/d, leg 5. **Male**, Pl. 54. 1a/b, habitus (lateral/dorsal); 1c, forehead (lateral); 1d, leg 2 basis; 1e/f, leg 3 basis. Pl. 55. 2a, urosome (dorsal); 2b, posterior prosome & urosome (lateral); 2c, leg 5 (posterolateral); 2d, right leg 5.

Razouls C. Desreumaux N., Kouwenberg J. and de Bovée F., 2005–2020. – Biodiversity of Marine Planktonic Copepods (morphology, geographical distribution and biological data). Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 03, 2021].