

Volume 30
Issue 5 Review-Special Issue (Part 2)

Article 41

Key to the species of Eucalanidae Pareucalanus occurring in the China Seas

Chang-tai Shih
National Taiwan Ocean University, ctshih@mail.ntou.edu.tw

Qing-Chao Chen
South China Sea Institute of Oceanology

Yang-Chi Lan
Fisheries Research Institute

Shih-Hui Hsiao
National Taipei University of Education

Chi-Yu Weng
Institute of Oceanography, National Taiwan University/Ocean Data Bank of the Ministry of National Science and Technology Council, Taiwan

Follow this and additional works at: <https://jmstt.ntou.edu.tw/journal>

 Part of the Fresh Water Studies Commons, Marine Biology Commons, Ocean Engineering Commons, Oceanography Commons, and the Other Oceanography and Atmospheric Sciences and Meteorology Commons

Recommended Citation

Shih, Chang-tai; Chen, Qing-Chao; Lan, Yang-Chi; Hsiao, Shih-Hui; and Weng, Chi-Yu (2022) "Key to the species of Eucalanidae Pareucalanus occurring in the China Seas," *Journal of Marine Science and Technology*. Vol. 30: Iss. 5, Article 41.

DOI: 10.51400/2709-6998.2632

Available at: <https://jmstt.ntou.edu.tw/journal/vol30/iss5/41>

This Review-Taxonomic Index is brought to you for free and open access by Journal of Marine Science and Technology. It has been accepted for inclusion in Journal of Marine Science and Technology by an authorized editor of Journal of Marine Science and Technology.

REVIEW-TAXONOMIC INDEX

Key to the Species of Eucalanidae *Pareucalanus* Occurring in the China Seas

Chang-tai Shih ^{a,*}, Qing-Chao Chen ^b, Yang-Chi Lan ^c, Shih-Hui Hsiao ^d, Chi-Yu Weng ^e

^a National Taiwan Ocean University, Taiwan

^b South China Sea Institute of Oceanology, China

^c Fisheries Research Institute, Taiwan

^d National Taipei University of Education, Taiwan

^e Institute of Oceanography, National Taiwan University/Ocean Data Bank of the Ministry of National Science and Technology Council, Taiwan

Pareucalanus attenuatus(3b/4a/f), *langae*(2a/4a/f), *sewelli*(3a/5b/f)

1a	Female	2
1b	Male	4
2a/1a	Cephalosome: width of shoulder less than 2 times width of triangular front. Mandible: basis with 4 setae..	<i>Pareucalanus langae</i>
2b	Cephalosome: width of shoulder more than 2 times width of triangular front. Mandible: basis with 2 setae	3
3a/2b	Antennule reaching distal border of prosome. Genital double-somite wider than long	<i>Pareucalanus sewelli</i>
3b	Antennule exceeding caudal rami by last 3 segments. Genital double-somite longer than wide	<i>Pareucalanus attenuatus</i>
4a/1b	Right leg 5 barely passing proximal border of left leg 5 segment 2	<i>Pareucalanus attenuatus</i>
4b	Right leg 5 nearly reaching midlength of left leg 5 segment 2.....	5
5a/4b	Right margin of usosomite 1 clearly convexed.....	<i>Pareucalanus langae</i>
5b	Right margin of urosomite 1 undulated	<i>Pareucalanus sewelli</i>


Available online 14 November 2022

* Corresponding author.

E-mail address: ctshih@mail.ntou.edu.tw (C.-t. Shih).



Pareucalanus attenuatus (Dana, 1849) (Size: female, 3.00–7.30 mm; male, 2.75–6.04 mm)




Pareucalanus attenuatus (Dana, 1849)

Phukham, 2008: Fig. 79. Female. a. habitus (dorsal); b. forehead (lateral); c. mandibular palp; d. mandibular cutting edge; e. posterior prosome and urosome (dorsal). Male. f. habitus (dorsal); g. posterior prosome & urosome (dorsal); h. leg 5 (right lateral).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed May 07, 2021].

Pareucalanus langae (Fleminger, 1973) (Size: female, 5.01–7.22 mm; male, 6.20–6.59 mm)




Pareucalanus langae (Fleminger, 1973)

Bradford-Grieve, 1994: Fig. 43. Female. A. habitus (dorsal/lateral); B. habitus (dorsal); C. urosome (dorsal); D. genital double-somite (lateral); E. mandibular palp. Male. F. urosome & leg 5 (right lateral).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en>. [Accessed Mayt 07, 2021].


Pareucalanus sewelli (Fleminger, 1973) (Size: female, 3.65–6.10 mm; male, 2.89–4.58 mm)



Pareucalanus sewelli (Fleminger, 1973)

Bradford-Grieve, 1994: Fig. 44. *Pareucalanus sewelli* (Fleminger, 1973), Female. A. habitus (dorsal and lateral); B. habitus (dorsal); C/D. urosome (dorsal/lateral); E. mandibular palp. Male. F. posterior prosome & urosome (dorsal). G. leg 5 (right lateral).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en> [Accessed May 07, 2021].

Pareucalanus sewelli (Fleminger, 1973)*Pareucalanus sewelli* (Fleminger, 1973)

Original. *Pareucalanus sewelli* (Fleminger, 1973), female, 4.60 mm, East of Taiwan (24°18'N, 121°46'E (Hualian, Heping), habitus (dorsal/lateral); rostrum; mandible.