



Journal of Marine Science and Technology



Volume 30
Issue 5 Review-Special Issue (Part 2)

Article 45

Key to the species of Heterorhabdidae *Heterorhabdus* occurring in the China Seas

Chang-tai Shih
National Taiwan Ocean University, ctshih@mail.ntou.edu.tw

Qing-Chao Chen
South China Sea Institute of Oceanology

Yang-Chi Lan
Fisheries Research Institute

Shih-Hui Hsiao
National Taipei University of Education

Chi-Yu Weng
Institute of Oceanography, National Taiwan University/Ocean Data Bank of the Ministry of National Science and Technology Council, Taiwan

Follow this and additional works at: <https://jmstt.ntou.edu.tw/journal>

 Part of the Fresh Water Studies Commons, Marine Biology Commons, Ocean Engineering Commons, Oceanography Commons, and the Other Oceanography and Atmospheric Sciences and Meteorology Commons

Recommended Citation

Shih, Chang-tai; Chen, Qing-Chao; Lan, Yang-Chi; Hsiao, Shih-Hui; and Weng, Chi-Yu (2022) "Key to the species of Heterorhabdidae *Heterorhabdus* occurring in the China Seas," *Journal of Marine Science and Technology*: Vol. 30: Iss. 5, Article 45.

DOI: 10.51400/2709-6998.2636

Available at: <https://jmstt.ntou.edu.tw/journal/vol30/iss5/45>

This Review-Taxonomic Index is brought to you for free and open access by Journal of Marine Science and Technology. It has been accepted for inclusion in Journal of Marine Science and Technology by an authorized editor of Journal of Marine Science and Technology.

REVIEW-TAXONOMIC INDEX

Key to the Species of Heterorhabdidae *Heterorhabdus* Occurring in the China Seas

Chang-tai Shih ^{a,*}, Qing-Chao Chen ^b, Yang-Chi Lan ^c, Shih-Hui Hsiao ^d, Chi-Yu Weng ^e

^a National Taiwan Ocean University, Taiwan

^b South China Sea Institute of Oceanology, China

^c Fisheries Research Institute, Taiwan

^d National Taipei University of Education, Taiwan

^e Institute of Oceanography, National Taiwan University/Ocean Data Bank of the Ministry of National Science and Technology Council, Taiwan

Heterorhabdus abyssalis(34a/f), *ankylocolus*(7a/26a/f), *clausi*(13b/28a/f), *cohibilis*(16b/34b/f), *confusibilis*(17a/35b/f), *egregious*(19b/37b/f), *fistulosus*(19a/37a/f), *habrosomus*(17b/32a/f), *insukae*(8a/23a/f), *oikoumenikis*(14/32b/f), *pacificus*(16a/35a/f), *papilliger*(18b/36a/f), *prolatus*(18a/36b/f), *quadrilobus*(7b/26b/f), *spinifrons*(8b/25a/f), *sub-spinifrons*(6a/24a/f), *tanner*(9a/30a/f)i, *tuberculus*(13a/28b/f) (*Spinifrons species group: *ankylocolus*, *insukae*, *quadrilobus*, *spinifrons*, *subspinifrons* (5); Papilliger species group: *papilliger*, *prolatus* (2); Fistulosus species group: *egregious*, *fistulosus* (2); Abyssalis species group: *abyssalis*, *clausi*, *cohibilis*, *confusibilis*, *habrosomus*, *oikoumenikis*, *pacificus*, *spinosus*, *tanneri*, *tuberculus* (10))

1a	Female	2
1b	Male	20
2a/1a	Maxilla: none of the spines on basal endite conspicuously serrated with long spinules. Maxilliped: middle coxal spine smoothly curved (spinifrons species group).....	5
2b	Maxilla: one of the spines on basal endite conspicuously serrated with long spinules. Maxilliped: middle coxal spine distinctly bent (73a)	3
3a/2b	Maxilla: unserrated spine on basal endite much shorter than serrated spine (abyssalis species group).....	9
3b	Maxilla: unserrated spine on basal endite nearly as long as or only a little shorter than serrated spine.....	4
4a/3b	In dorsal view, midanterior tubercular process of cephalosome pointed. Maxilliped: basis without large elevated pores (papilliger species group)	18
4b	In dorsal view, midanterior tubercular process of cephalosome somewhat truncate. Maxilliped: basis with large elevated pore on medial margin close to distal end (fistulosus species group).....	19
5a/2b	Anterior wall of genital prominence with a pair of rounded lobes.....	6
5b	Anterior wall of genital prominence without such lobes.....	8
6a/5a	Genital prominence reaching posterior end of genital double-somite. Maxillule: endopod with 1 seta	<i>Heterorhabdus subspinifrons</i>
6b	Genital prominence short of reaching posterior end of genital double-somite. Maxillule: endopod with 3 setae	7
7a/6b	Genital prominence without posterior lobes	<i>Heterorhabdus ankylocolus</i>
7b	Genital prominence with posterior lobes	<i>Heterorhabdus quadrilobus</i>
8a/5b	Genital-double somite in dorsal view relatively long (1.4 time combined length of urosomites 2 & 3), with wavy lateral margins. In lateral view, genital prominence not reaching posterior end of genital double-somite. Maxillule without seta on basal epipodite and 1 seta on endopod.....	<i>Heterorhabdus insukae</i>
8b	Genital double-somite in dorsal view relatively short (slightly shorter than combined length of urosomites 2 and 3), with smoothly bulging lateral margins. In lateral view, genital prominence reaching posterior end of genital double-somite, without posterior lobe. Maxillule with 1 seta on basal epipodite, 3 setae on endopod.....	<i>Heterorhabdus spinifrons</i>
9a/3a	In lateral view, genital double-somite with a tubercular outgrowth posteriorly on ventral margin. In lateral view, genital operculum short of reaching posterior end of genital double-somite	<i>Heterorhabdus tanneri</i>
9b	In lateral view, genital double-somite without such tubercular outgrowth.....	10

Available online 14 November 2022

* Corresponding author.

E-mail address: ctshih@mail.ntou.edu.tw (C.-t. Shih).



10a/9b	Laterally, genital operculum reaching close to posterior end of genital double-somite.....	11
10b	Laterally, genital operculum far short of reaching posterior end of genital double-somite	12
11a/10a	In lateral view, genital flange short and high, distinctly separate from posterior ventral margin of genital double-somite; Maxilla: posterior subterminal spine of distal coxal endite about 1/5 length of 2nd saberlike spine ... <i>Heterorhabdus spinosus</i>	
11b	In lateral view, genital flange elongate and low, separate from posterior ventral margin of genital double-somite by a shallow notch; Maxilla: posterior subterminal spine of distal coxal endite almost 1/2 length of 2nd saberlike spine.....	<i>Heterorhabdus abyssalis</i>
12a/10b	In lateral view, right and left genital flanges asymmetrical	13
12b	In lateral view, right and left genital flanges symmetrical	14
13a/10b	In lateral view, left genital flange high, like a tubercular outgrowth.....	<i>Heterorhabdus tuberculatus</i>
13b	In lateral view, left genital flange low, smoothly arched. Laterally genital double-somite relatively short, typically with a length-depth ratio of 100:89	<i>Heterorhabdus clausi</i>
14a/12b	In lateral view, genital prominence gradually merging onto ventral wall of genital double-somite. In dorsal view, genital double-somite slightly asymmetrical.....	<i>Heterorhabdus oikoumenikis</i>
14b	In lateral view, genital prominence distinctly separate from ventral wall of genital double-somite	15
15a/14b	In dorsal and ventral views, lateral swelling of genital double-somite reaching posterior end of the somite.....	16
15b	In dorsal and ventral views, lateral swelling of genital double-somite short of reaching posterior end of the somite	17
16a/15a	In lateral view, genital prominence extending close to posterior end of genital double-somite; the somite with a conspicuous dorsal hump	<i>Heterorhabdus pacificus</i>
16b	In lateral view, genital prominence far short of extending to posterior end of genital double-somite; the somite with a low dorsal hump. In lateral view, genital operculum far short of reaching posterior end of genital prominence	<i>Heterorhabdus cohibilis</i>
17a/15b	In lateral view, genital prominence reaching close to posterior end of genital double-somite. Maxilla: posterior subterminal spine of distal coxal endite shorter than one-half length of saberlike spine.....	<i>Heterorhabdus confusibilis</i>
17b	In lateral view, genital prominence far short of reaching posterior end of genital double-somite. Maxilla: posterior sub-terminal spine of distal coxal endite longer than one-half length of saberlike spine	<i>Heterorhabdus habrosomus</i>
18a/4a	In lateral view, genital prominence extending far short to posterior end of genital double-somite	<i>Heterorhabdus prolatus</i>
18b	In lateral view, genital prominence extending down to posterior end of genital double-somite.....	<i>Heterorhabdus papilliger</i>
19a/4b	In dorsal view, genital double-somite widest in anterior half; in lateral view, genital prominence far short of reaching posterior end of genital double-somite	<i>Heterorhabdus fistulosus</i>
19b	In dorsal view, genital double-somite widest in posterior half; in lateral view, genital prominence extending down to posterior end of genital double-somite	<i>Heterorhabdus egregius</i>
20a/1a	Maxilla: none of the spines on basal endite conspicuously serrated with long spinules. Maxilliped: middle coxal spine smoothly curved (<i>spinifrons</i> species group).....	23
20b	Maxilla: one of the spines on basal endite conspicuously serrated with long spinules. Maxilliped: middle coxal spine distinctly bent	21
21a/2b	Maxilla: unserrated spine on basal endite much shorter than serrated spine (<i>abyssalis</i> species group).....	27
21b	Maxilla: unserrated spine on basal endite nearly as long as or only a little shorter than serrated spine	22
22a/3b	In dorsal view, midanterior tubercular process of cephalosome pointed. Maxilliped: basis without large elevated pores (<i>papilliger</i> species group)	36
22b	In dorsal view, midanterior tubercular process of cephalosome somewhat truncate. Maxilliped: basis with large elevated pore on medial margin close to distal end (<i>fistulosus</i> species group)	37
23a/20a	Right leg 5: basal inner lobe relatively short and conical; medial projection of exopod segment 2 with large terminal spiniform process and poorly developed distal lobe	<i>Heterorhabdus insukae</i>
23b	Right leg 5: basal inner lobe relatively long and somewhat finger-like; medial projection of exopod segment 2 with small terminal spiniform process and well developed distal lobe.....	24
24a/23b	Right leg 5: basal inner lobe almost as long as basis, pointing distally. Maxillule: endopod with 1 seta	<i>Heterorhabdus subspinifrons</i>
24b	Right leg 5: basal inner lobe much shorter than basis, pointing distomedially. Maxillule: endopod with 3 setae	25
25a/24b	Left leg 5: exopod segment 2 with outer spine longer than conical process; right leg 5 exopod segment 3: terminal spine about 1/6 length of the segment.....	<i>Heterorhabdus spinifrons</i>
25b	Left leg 5: exopod segment 2 with outer spine equal to or shorter than conical process; right leg 5 exopod segment 3: terminal spine about 1/3 length of the segment	26
26a/25b	Right leg 5 exopod segment 3: lateral margin strongly bent at about 1/5 length of the segment from distal end; left leg 5 exopod segment 1 with a straight inner margin	<i>Heterorhabdus ankylocolus</i>
26b	Right leg 5 exopod segment 3: lateral margin slightly or moderately bent at about 1/5 length of the segment from distal end; left leg 5 exopod segment 1 with a semicircular bulging at distal inner margin	<i>Heterorhabdus quadrilobus</i>
27a/21a	Left leg 5: basis with well-developed inner lobe	28
27b	Left leg 5: basis without well-developed inner lobe	29
28a/27a	Left leg 5: basal inner lobe wide, width 2.5 times length	<i>Heterorhabdus clausi</i>
28b	Left leg 5: basal inner lobe long, length 1.5 times width	<i>Heterorhabdus tuberculatus</i>
29a/27b	Right leg 5: basal inner lobe armed with thick lancet-like bristles	<i>Heterorhabdus spinosus</i>
29b	Right leg 5: basal inner lobe armed with normal bristles.....	30
30a/29b	Right leg 5: basal inner lobe arising from anterior surface of the segment	<i>Heterorhabdus tanneri</i>
30b	Right leg 5: basal inner lobe arising from medial or anteromedial surface of the segment.....	31

31a/30b	Left leg 5 exopod segment 2: outer spine arising from a conical process	32
31b	Left leg 5 exopod segment 2: outer spine not arising from a conical process	33
32a/31a	Right leg 5: basal inner lobe sigmoid-shaped	<i>Heterorhabdus habrosomus</i>
32b	Right leg 5: basal inner lobe straight.....	<i>Heterorhabdus oikoumenikis</i>
33a/31b	Right leg 5: anteriorly and when tilted clockwise, basal inner lobe recurved by folding	34
33b	Right leg 5: anteriorly and when tilted clockwise, basal inner lobe recurved without folding.....	35
34a/33a	Right leg 5: anteriorly, basal inner lobe strongly folded. Left leg 5 exopod segment 3: medial spine 2 times as long as lateral spine.....	<i>Heterorhabdus abyssalis</i>
34b	Right leg 5: anteriorly, basal inner lobe slightly folded. Left leg 5 exopod segment 3: medial spine nearly 3 times as long as lateral spine.....	<i>Heterorhabdus cohibilis</i>
35a/33b	Right leg 5: lateral margin of basal inner lobe forming a narrow, rounded arch with medial margin of the segment. Left leg 5 exopod segment 3: medial spine nearly 3 times as long as lateral spine	<i>Heterorhabdus pacificus</i>
35b	Right leg 5: lateral margin of basal inner lobe forming a narrow angle with medial margin of the segment. Left leg 5 exopod segment 3 less than 2 times as long as lateral spine	<i>Heterorhabdus confusibilis</i>
36a/22a	Right leg 5: basal inner lobe slightly shorter than 1/2 length of the segment and pointing distally; medial projection of exopod segment 2 with a large, rounded, plumose proximal lobe and without a definable distal lobe	<i>Heterorhabdus papillige</i>
36b	Right leg 5: basal inner lobe elongate, about 2/3 length of the segment and pointing distomedially; medial projection of exopod segment 2 elongate, without terminal enlargement and with nearly parallel proximal and distal sides	<i>Heterorhabdus prolatus</i>
37a/22b	Right leg 5: basal inner lobe elongate; left leg 5 exopod segment 2: lateral conical process and outer spine elongate, with their combined length about two-thirds length of the segment	<i>Heterorhabdus fistulosus</i>
37b	Right leg 5: basal inner lobe wide; left leg 5 exopod segment 2: lateral conical process and outer spine short, with their combined length about 1/4 length of the segment.....	<i>Heterorhabdus egregius</i>

Heterorhabdus abyssalis (Giesbrecht, 1889) (Size: female, 2.09–3.73 mm; male, 2.00–3.40 mm)




Fig. 106





Fig. 107

Heterorhabdus abyssalis (Giesbrecht, 1889)

Park, 2000: Fig. 106. Female. a. habitus (lateral); b/c, urosome (lateral/dorsal); d, genital double-somite (left); e/f/g, genital double-somite with operculum open (right/left/ventral); h. maxilla, distal coxal endite; i, leg 1 exopod. Fig. 107. Male. a. leg 5 (anterior); b. right leg 5, distal end of exopod (lateral); c/d/e. right leg 5 basis (anterior/anterior/posterior); f. left leg 5 exopod (anterior); g/h right leg 5 exopod segment 2 (anterior/posterior).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en> [Accessed May 10, 2021].

Heterorhabdus ankylocolus Park, 2000 (Size: female, 2.80–3.24 mm; male: 2.72–2.92 mm)



Heterorhabdus ankylocolus Park, 2000

Park, 2000: Fig. 67. Female: a. urosome (lateral); b/c/d. genital double-somite (dorsal/left/ventral). Male. e, left leg 5 (anterior, endopod omitted); f. left leg 5, exopod (anterior); g. right leg 5 (anterior, distal exopodal segments omitted); h. right leg 5 (posterior, endopod omitted); i. right leg 5 exopod (anterior); j. right leg 5, distal end of exopod (lateral).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en> [Accessed May 10, 2021].

Heterorhabdus caribbeanensis Park, 1970 (Size:female, 1.74–2.52 mm; male, 1.92–2.40 mm)




Fig. 68

Heterorhabdus caribbeanensis Park, 1970




Fig. 69

T. Park, 2000, Fig. 68. Female: a, habitus (left side); b, urosome (dorsal); c/d, forehead (dorsal/Lateral); e/f, genital double-somite (lateral/ventral); g, left Maxillule (posterior) Fig. 69. Female: a, left maxilla (posterior); b, right maxilliped (anterior). Male: c, leg 5 (distal exopodal segments omitted), anterior; d, right leg 5 (endopod omitted), posterior; e, exopod of left P5 (anterior); f, first 2 exopodal segments of left P5 (anterior). Female: a, left maxilla (posterior); b, right maxilliped (anterior). Male: c, leg 5 (distal exopodal segments omitted), anterior; d, right leg 5 (endopod omitted), posterior; e, exopod of left leg 5 (anterior); f, first 2 exopodal segments of left leg 5 (anterior)

Heterorhabdus clausi (Giesbrecht, 1889) (Size: female, 2.00–2.70 mm; male: 2.00–2.50 mm)




Fig. 88

Heterorhabdus clausi (Giesbrecht, 1889)





Fig. 89

Park, 2000: Fig. 88. Female. a, habitus (lateral); b/c, urosome (lateral/dorsal). d/e, genital double-somite (left/right). gfl = genital flange. Fig. 89. Female: a, genital somite with operculum open (left side); b, genital somite (ventral); c, genital somite with operculum open (right). Male: d, leg 5 (anterior); e, basipod of left leg 5 (anterior).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en> [Accessed May 07, 2021].

Heterorhabdus cohibilis Park, 2000 (Size: female, 2.80–3.60 mm; male: 2.68–3.32 mm)




Heterorhabdus cohibilis Park, 2000


Park, 2000: Fig. 99. *Heterorhabdus cohibilis* Park, 2000, Female. a. habitus (lateral); b. urosome (dorsal); c/d. genital double-somite (ventral/lateral). Male. e. right leg 5 (anterior); f/g. right leg 5 basis (anterior); h. left leg 5 (anterior); i. left leg 5 exopod (anterior).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en> [Accessed May 8, 2021].

Heterorhabdus confusibilis Park, 2000 (Size: female, 2.52–3.44 mm; male, 2.56–3.08 mm)



Female



Male

Heterorhabdus confusibilis Park, 2000

Park, 2000: Fig. 110. *Heterorhabdus confusibilis* Park, 2000, Female. a. habitus (lateral); b. urosome (dorsal); c/d. forehead (lateral/dorsal); e. genital double-somite (lateral); f. genital double-somite (with operculum open); g. genital double-somite (ventral); h. left maxilla (posterior); i. leg 5 (anterior). Fig. 111. Male. a. leg 5 (anterior); b. left leg 5 (anterior, endopod omitted); c. left leg 5 exopod (anterior); d. right leg 5 (posterior, endopod omitted); e. right leg 5, distal end of exopod (lateral); f/g/h. right leg 5, basis (anterior/anterior/posterior); i/j. right leg 5, exopod segment 2 (anterior/posterior).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en> [Accessed May 10, 2021].

Heterorhabdus egregius Heptner, 1972 (Size: female, 2.24–3.40 mm; male: 2.28–3.70 mm)




Fig. 82

Heterorhabdus egregius Heptner, 1972





Fig. 83

Park, 2000: Fig. 82. *Heterorhabdus egregius* Heptner, 1972, Female. a. habitus (lateral); b. urosome (dorsal); c/d, forehead (lateral/dorsal); e/f. genital double-somite (lateral/ventral); g. left maxilla (posterior); h. right maxilliped (anterior); i/j/k. right maxilliped, basis (anterior, from 3 different specimens). Fig. 83. Female. a. leg 5 (anterior); b. leg 5 exopod (posterior). Male. c. leg 5 (anterior); d/e. right leg 5, basis (anterior/posterior); f. left leg 5, basis (anterior); g/h. right leg 5 exopod segment 2 (anterior/posterior); i. left leg 5 exopod (anterior)

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en> [Accessed May 10, 2021].

Heterorhabdus egregius Heptner, 1972*Heterorhabdus egregius* Heptner, 1972

Original. *Heterorhabdus egregius* Heptner, 1972, female, 2.86 mm, east of Taiwan, habitus (dorsal/lateral), maxilliped.

Heterorhabdus fistulosus Tanaka, 1964 (Size: female, 3.39–3.95; male, 3.40–3.85 mm)




Fig. 80

Heterorhabdus fistulosus Tanaka, 1964




Fig. 81

Park, 2000: Fig. 80. Female. a. habitus (lateral); b/c. urosome (lateral/dorsal); d. genital double-somite (lateral); e. genital douoble-somite with opercululum open (lateral); f. genital double-somite (ventral); g/h/i. genital douoble-somite with a tubercular ouotgrowth (dorsal/left/right); j. left maxilla (posterior). Fig. 81. Female: a, right maxilliped (anterior); b/c, distal part of basis of right maxilliped (anterior, from 2 specimens); d, leg 1 exopod (anterior); e, leg 5 exopod.(posterior). Male: f, leg 5 (anterior); g/h, right leg 5 exopod segment 2 (anterior/posterior); i, left leg 5 exopod (anterior).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en> [Accessed May 08, 2021].

Heterorhabdus habrosomus Parak, 2000 (Size: female, 2.16–2.56; male, 2.14–3.08 mm)




Fig. 84

Heterorhabdus habrosomus Parak, 2000




Fig. 85

Park, 2000: Fig. 84. As *Heterorhabdus egregius* Heptner, 1972, Female. a. habitus (lateral); b/c. urosome (lateral/dorsal); d/e. forehead (lateral/dorsal); f/g. genital double-somite (lateral, from 2 specimens); h. genital double-somite (ventral); i. left maxilla (posterior); j. right maxilliped (anterior). Fig. 85. Female. a/b/c/d/e. leg 1/2/3/4/5 exopod (anterior); f. leg 5 exopod (posterior). Male. g. leg 5 (anterior); h. left leg 5 basis (anterior); i. left leg 5 exopod (anterior); j. right leg 5 (posterior, endopod omitted); k. right leg 5 (anterior, distal exopod segments omitted).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en> [Accessed May 07, 2021].

Heterorhabdus insukae Park, 2000 (Size: female, 1.78–2.56; male, 2.12–2.48 mm)




Fig. 70

Heterorhabdus insukae Park, 2000




Fig. 71

Park, 2000: Fig. 70. *Heterorhabdus insukae* Park, 2000, Female. a. habitus (lateral); b/c. urosome (lateral/dorsal); d/e, forehead (lateral/dorsal); f/g. genital double-somite (lateral/ventral); h. left maxillule (posterior); i. left maxilla (posterior). Fig. 71. Female. a. right maxilliped (anterior); b. leg 5 (posterior). Male. c. leg 5 (anterior); d. right leg 5 basis (anterior); e. right leg 5 exopod (posterior); f. right leg 5, exopod segment 2 (anterior); g. left leg 5 (anterior, endopod omitted); h. left leg 5, exopod segments 1–2 (anterior).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en> [Accessed May 08, 2021].

Heterorhabdus oikoumenikis Park, 2000 (Size: female, 2.60–3.52; male, 2.40–3.28 mm)




Fig. 94

Heterorhabdus oikoumenikis Park, 2000




Fig. 95

Park, 2000: Fig. 94. *Heterorhabdus oikoumenikis* Park, 2000, Female. a. habitus (lateral); b/c. urosome (lateral/dorsal); d/e. forehead (lateral/dorsal); f/g. genital double-somite (ventral/lateral); h. genital double-somite (lateral, from another specimen); i. left maxilla (posterior). Fig. 95. Female. a/b. leg 5 (anterior/posterior). Male. c. leg 5 (anterior); d. right leg 5 exopod (anterior); e. right leg 5, exopod segment 2; f/g. right leg 5, basis (anterior/posterior); h/i. right leg 5, basis (anterior/posterior, from another specimen); j. left leg 5, exopod (anterior); k. right leg 5, distal end of exopod (lateral).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en> [Accessed May 07, 2021].

Heterorhabdus pacificus Brodsky, 1950 (Size: female, 1.92–3.84; male, 2.90–3.96 mm)




Fig. 108

Heterorhabdus pacificus Brodsky, 1950





Fig. 109


Park, 2000: Fig. 108. *Heterorhabdus pacificus* Brodsky, 1950, Female. a. habitus (lateral); b. urosome (dorsal); c/d. forehead (lateral/dorsal); e/f. genital double-somite (ventral/lateral); g. left maxilla (posterior); h. leg 5 (anterior); i. leg 5 exopod (posterior). Fig. 109. Male. a. leg 5 (anterior); b/c. right leg 5 (anterior/posterior, distal segments of rami omitted); d. right leg 5, basis (anterior); e. right leg 5, distal end of exopod (lateral); f. left leg 5 exopod (anterior); g/h. right leg 5, exopod segment 2 (anterior/posterior).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en> [Accessed May 07, 2021].


Heterorhabdus papilliger Claus, 1863 (Size: female, 1.60–2.66; male, 1.60–2.65 mm)



Grice, 1962



Chen & Zhang, 1965



Heterorhabdus papilliger Claus, 1863

Grice, 1962: pl. 24. *Heterorhabdus papilliger* Claus, 1863, Female. 12. habitus (lateral)

Chen & Zhang, 1965: pl. 34. *Heterorhabdus papilliger* Claus, 1863, Female. 1/2. habitus (dorsal/lateral); 3. maxilliped; 4. right leg 5 (anterior). Male. 5. habitus (dorsal); 6. leg 5 (anterior).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en> [Accessed May 07, 2021].

Heterorhabdus prolatus Park, 2000 (Size: female, 1.68–2.14; male, 1.36–2.14 mm)




Fig. 78




Fig. 79

Heterorhabdus prolatus Park, 2000

Park, 2000: Fig. 78. *Heterorhabdus prolatus* Park, 2000, Female. a. habitus (lateral); b/c. urosome (lateral/dorsal); d/e. forehead (lateral/dorsal); f/g/h. genital double-somite (left/ventral/right). Fig. 79. Female. a. left maxilla (posterior); b. leg 5 (anterior). Male. c. right leg 5 (anterior, endopod omitted); d. left leg 5 (anterior); e. leg 5 exopod (anterior).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en> [Accessed May 05, 2021].

Heterorhabdus quadrilobus Park, 2000 (Size: female, 2.88–3.84; male, 2.48–3.68 mm)




Fig. 65

Heterorhabdus quadrilobus Park, 2000




Fig. 66

Park, 2000: Fig. 65. *Heterorhabdus quadrilobus* Park, 2000, Female. a/b. urosome (lateral/dorsal); c/d/e. genital double-somite (left/right/ventral). Fig. 66. Female. a. right maxilliped (anterior). Male. b. leg 5 (anterior); c. right leg 5, basis (anterior); d. left leg 5 (anterior, endopod omitted); e. right leg 5 exopod (posterior); f. right leg 5, distal end of exopod (lateral).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en> [Accessed May 07, 2021].

Heterorhabdus spinifrons Claus, 1863 (Size: female, 2.86–4.20; male, 2.20–3.84 mm)




Fig. 59




Fig. 60
Heterorhabdus spinifrons Claus, 1863





Fig. 61

Park, 2000: Fig. 59. Female. a. habitus (lateral); b/c. urosome (lateral/dorsal); d/e. forehead (lateral/dorsal); f/g/h/i. genital double-somite (dorsal/right/left/ventral); j. left antennule (ventral). Fig. 60. Female. a. left antenna (posterior); b. left antenna, endopod segment 1 (anterior); c. left mandible (posterior); d. left mandible gnathobase (posterior); e. right mandible, cutting edge (posterior); f. left maxillule (posterior); g. left maxilla (posterior); h. right maxilliped (anterior); i. leg 1; j. let 1, basis; k/l/m. leg 2/3/4 (anterior). Fig. 61. Female. a. leg 5 (anterior). Male. b. urosome (dorsal); c. left antennule (dorsal); d. leg 5 (anterior); e. right leg 5 (anterior, endopod omitted); f. right leg 5, distal end of exopod (lateral); g. right leg 5, exopod (posterior); h. right leg 5 exopod segment 2 (anterior); i. left leg 5, distal exopod segments (anterior).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en> [Accessed May 07, 2021].

Heterorhabdus subspinifrons Tanaka, 1964 (Size: female, 2.27–3.20; male, 2.11–3.16 mm)



Heterorhabdus subspinifrons Tanaka, 1964

Park, 2000: Fig. 64. *Heterorhabdus subspinifrons* Tanaka, 1964, Female. a. urosome (dorsal); b/c. genital double-somite (lateral/ventral); d. left maxillule (posterior); e. left maxilla, distal end (posterior); f. right maxilliped (anterior); g. leg 5 (anterior). Male. h. right leg 5 (anterior, distal exopod segments omitted); i. right leg 5 (posterior, endopod omitted); j. right leg 5, distal end of exopod (lateral); k. left leg 5 (anterior, endopod omitted).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en> [Accessed May 07, 2021].

Heterorhabdus tanneri Giesbrecht, 1895 (Size: female, 3.08–4.90; male, 3.00–4.25 mm)




Fig. 112




Fig. 113

Heterorhabdus tanneri Giesbrecht, 1895

Park, 2000: Fig. 112. *Heterorhabdus tanneri* Giesbrecht, 1895, Female. a. habitus (lateral); b. urosome (dorsal); c/d. genital double-somite (ventral/lateral); e. genital double-somite with operculum partially open (lateral); f. left maxilla (posterior); g. leg 5 (posterior). Fig. 113. Male. a. leg 5 (anterior); b. right leg 5, distal end of exopod (lateral); c. left leg 5, basis (anterior); d/e. right leg 5 (posterior/anterior, endopod omitted); f. left leg 5, exopod (anterior).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en> [Accessed May 07, 2021].

Heterorhabdus tuberculatus Park, 2000 (Size: female, 2.24–2.84; male, 2.24–2.68 mm)




Fig. 90




Fig. 91

Heterorhabdus tuberculatus Park, 2000

Park, 2000: Fig. 90. *Heterorhabdus tuberculatus* Park, 2000, Female. a. habitus (lateral); b/c. urosome (dorsal/lateral); d/e/f. genital double-somite (ventral/left/right); g. genital double-somite (lateral); h. genital double-somite with operculum open (lateral). Fig. 91. Female. a. genital double-somite (dorsal); b. genital double-somite from another specimen (dorsal); c/d. abnormal genital double-somite (lateral). Male. e. leg 5 (anterior); f. leg 5 (anterior, distal segments omitted).

Adapted from Razouls C., de Bovée F., Kouwenberg J. and Desreumaux N., 2005–2020. Diversity and Geographic Distribution of Marine Planktonic Copepods. Sorbonne University, CNRS. Available at <http://copepodes.obs-banyuls.fr/en> [Accessed May 07, 2021].